



Complete Genome Sequence of Avian Coronavirus Strain GA08 (GI-27 Lineage)

 Iryna V. Goraichuk,^a James F. Davis,^b Arun B. Kulkarni,^b  Claudio L. Afonso,^a  David L. Suarez^a

^aExotic and Emerging Avian Viral Disease Research Unit, Southeast Poultry Research Laboratory, U.S. National Poultry Research Center, ARS, USDA, Athens, Georgia, USA

^bGeorgia Poultry Laboratory Network, Gainesville, Georgia, USA

ABSTRACT *Avian coronavirus*, also known as infectious bronchitis virus, is a highly contagious respiratory pathogen of chickens that is responsible for major economic losses to the poultry industry around the globe. Here, we report the complete genome sequence of strain GA08 of the GI-27 lineage, isolated from a fecal sample from a broiler chicken collected in Georgia in 2015.

A *avian coronavirus*, commonly known as infectious bronchitis virus (IBV), is a member of the genus *Gammacoronavirus*, family *Coronaviridae*, order *Nidovirales* (1–3). Since the first identification of IBV in the 1930s in the United States, more than 20 different serotypes and variants have been identified worldwide (4, 5). In 2007 and 2008, an outbreak of bronchitis in broilers in North Georgia led to the identification of a new IBV strain designated GA08, which was further classified into the separate lineage GI-27 (6, 7). The GA08 IBV strain has since been isolated on farms in Alabama, Tennessee, and Kentucky (8; <https://quickvet.net/news-and-media/zoetis-receives-usda-conditional-license-for-first-vaccine-to-control-georgia-08-strain-of-infectious-bronchitis-virus-in-poultr.aspx>). Over the past years, its incidence and severity have increased dramatically, resulting in significant losses. To date, only partial genomes of the GA08 strain viruses are available in public databases (6, 7). In this study, we report the complete sequence of the IBV GA08 strain.

A fecal sample from an 11-day-old broiler chicken was collected at a commercial farm in Georgia in 2015. Feces were passed sequentially through 1.2- μ m- and 0.45- μ m-pore-size filters (Merck Millipore, USA) to remove bacteria and particles with large cell sizes. Total nucleic acids were isolated from the filtered lysate using the DNeasy blood and tissue kit (Qiagen, Germany), followed by DNase treatment with the Turbo DNA-free kit (Ambion, USA) to remove host DNA according to the manufacturer's recommendations. Sequence-independent single-primer amplification (9) was used to produce random amplicons that were processed using the Nextera XT DNA library preparation kit (Illumina, USA). The distribution size and concentration of the prepared library were checked on a Bioanalyzer 2100 instrument, using the Agilent high-sensitivity (HS) DNA kit (Agilent Technologies, Germany), and on a Qubit fluorometer using the double-stranded DNA (dsDNA) HS assay kit (Life Technologies, USA), respectively. Next-generation paired-end sequencing (2 \times 250 bp) was performed on an Illumina MiSeq instrument using the 500-cycle MiSeq reagent kit v2 (Illumina, USA). Sequence data were assembled using a *de novo* approach and utilizing MIRA3 version 0.0.1 (10) within a customized workflow on the Galaxy platform (11), as described previously (12, 13). The final outputs of the analysis workflow included a set of summary statistics on the run and the assembly. A total of 2,424,105 raw paired-end reads with a mean length of 174 bp were generated. The genome consensus of the isolate, designated GA/1476/2015, was called from 36,893 IBV reads using BWA-MEM (14). This resulted in a median read depth coverage of 141

Citation Goraichuk IV, Davis JF, Kulkarni AB, Afonso CL, Suarez DL. 2020. Complete genome sequence of *Avian coronavirus* strain GA08 (GI-27 lineage). *Microbiol Resour Announc* 9:e00068-20. <https://doi.org/10.1128/MRA.00068-20>.

Editor Simon Roux, DOE Joint Genome Institute

This is a work of the U.S. Government and is not subject to copyright protection in the United States. Foreign copyrights may apply.

Address correspondence to David L. Suarez, david.suarez@usda.gov.

Received 21 January 2020

Accepted 3 February 2020

Published 27 February 2020




FIG 1 Phylogenetic analysis of IBV isolates of the GA08 strain based on the complete S1 gene sequences. The S1 gene sequences of 21 IBV isolates were downloaded from the NCBI GenBank database. Together with the sequence obtained in the current study, all sequences were subjected to multiple alignment using the ClustalW algorithm. The phylogenetic tree was constructed using the maximum likelihood method based on the general time-reversible model in MEGA version 7.0.26. The tree with the highest log likelihood ($-6,129.42$) is shown. The percentage of replicate trees in which the associated taxa clustered together in the bootstrap test (1,000 replicates) is shown next to each branch. The tree is drawn to scale, with branch lengths measured in the number of substitutions per site. The analysis involved 22 nucleotide sequences (a sequence from the GI-9 lineage is included as an outgroup). All positions containing gaps and missing data were eliminated. There was a total of 1,608 positions in the final data set. The isolates used in this study are shown in red.

reads. The final genome consensus was 27,650 nucleotides long, excluding the poly(A) tail (100% genome coverage), and had a GC content of 38.1%. The open reading frames (ORFs) were identified using Geneious version 11.1.5 and confirmed by alignment with published IBV genomes. The genome has the typical genetic structure of all IBV strains and contains 13 ORFs (5'-1a/1b-S-3a-3b-E-M-4b-4c-5a-5b-N-6b-3'). A preliminary BLAST comparison of the GA/1476/2015 genome to currently available full-length IBV genome sequences showed the highest (95.31%) nucleotide identity to the Arkansas strain ArkDPI11 (GenBank accession number [EU418976](https://www.ncbi.nlm.nih.gov/nuccore/EU418976)) belonging to the GI-9 lineage (15, 16). Detailed phylogenetic analysis based on the complete coding sequence of the S1 gene (8) revealed that the GA/1476/2015 isolate is a member of the GI-27 lineage and forms a distinct cluster along with the lineage prototype strain Georgia 08/2008 (99.88% nucleotide identity; GenBank accession number [GU301925](https://www.ncbi.nlm.nih.gov/nuccore/GU301925)) (6) (Fig. 1). Currently, there are no IBV reference full-genome sequences available for the GA08 strain isolates, only sequences of the S1 gene (6, 7). This complete genome sequence information of an IBV GA08 strain isolate will be useful for in-depth understanding of IBV evolution, as well as facilitate future development of new vaccines.

Data availability. The complete genome sequence of the *Avian coronavirus* GA/1476/2015 isolate of the GA08 strain has been deposited in GenBank under accession number [MN599049](https://www.ncbi.nlm.nih.gov/nuccore/MN599049). Raw data were deposited under SRA accession number [SRR10742650](https://www.ncbi.nlm.nih.gov/sra/SRR10742650), BioSample number [SAMN13088878](https://www.ncbi.nlm.nih.gov/biosample/SAMN13088878), and BioProject number [PRJNA556282](https://www.ncbi.nlm.nih.gov/bioproject/PRJNA556282).

ACKNOWLEDGMENTS

The mention of trade names or commercial products in this publication is solely for the purpose of providing specific information and does not imply recommendation or endorsement by the U.S. Department of Agriculture.

This study was supported by USDA CRIS project 6040-32000-072.

REFERENCES

- Jackwood MW, de Wit JJ. 2013. Infectious bronchitis, p 139–159. *In* Swayne DEGJ, McDougald LR, Nolan LK, Suarez DL, Nair V (ed), *Diseases of poultry*, 13th ed. John Wiley and Sons, Inc., Hoboken, NJ.
- Cavanagh D. 2007. Coronavirus avian infectious bronchitis virus. *Vet Res* 38:281–297. <https://doi.org/10.1051/vetres:2006055>.
- King AMQ, Lefkowitz EJ, Mushegian AR, Adams MJ, Dutilh BE, Gorbalenya AE, Harrach B, Harrison RL, Junglen S, Knowles NJ, Kropinski AM, Krupovic M, Kuhn JH, Nibert ML, Rubino L, Sabanadzovic S, Sanfacon H, Siddell SG, Simmonds P, Varsani A, Zerbini FM, Davison AJ. 2018. Changes to taxonomy and the International Code of Virus Classification and Nomenclature ratified by the International Committee on Taxonomy of Viruses (2018). *Arch Virol* 163:2601–2631. <https://doi.org/10.1007/s00705-018-3847-1>.
- Roh HJ, Hilt DA, Jackwood MW. 2013. Simultaneous detection of five major serotypes of Avian coronavirus by a multiplex microspheres-based assay. *J Vet Diagn Invest* 25:458–466. <https://doi.org/10.1177/1040638713493906>.
- Liu S, Chen J, Han Z, Zhang Q, Shao Y, Kong X, Tong G. 2006. Infectious bronchitis virus: S1 gene characteristics of vaccines used in China and efficacy of vaccination against heterologous strains from China. *Avian Pathol* 35:394–399. <https://doi.org/10.1080/03079450600920984>.
- Jackwood MW, Hilt DA, Sellers HS, Williams SM, Lasher HN. 2010. Rapid heat-treatment attenuation of infectious bronchitis virus. *Avian Pathol* 39:227–233. <https://doi.org/10.1080/03079451003801516>.
- Kulkarni AB, Resurreccion RS. 2010. Genotyping of newly isolated infectious bronchitis virus isolates from northeastern Georgia. *Avian Dis* 54:1144–1151. <https://doi.org/10.1637/9358-040510-Reg.1>.
- Valastro V, Holmes EC, Britton P, Fusaro A, Jackwood MW, Cattoli G, Monne I. 2016. S1 gene-based phylogeny of infectious bronchitis virus: an attempt to harmonize virus classification. *Infect Genet Evol* 39:349–364. <https://doi.org/10.1016/j.meegid.2016.02.015>.
- Chrzastek K, Lee DH, Smith D, Sharma P, Suarez DL, Pantin-Jackwood M, Kapczynski DR. 2017. Use of sequence-independent, single-primer-amplification (SISPA) for rapid detection, identification, and characterization of avian RNA viruses. *Virology* 509:159–166. <https://doi.org/10.1016/j.virol.2017.06.019>.
- Chevreur B, Wetter T, Suhai S. 1999. Genome sequence assembly using trace signals and additional sequence information, p 45–56. *In* *Computer science and biology: proceeding of the German Conference on Bioinformatics*, GCB '99. GCB, Hanover, Germany.
- Afgan E, Baker D, van den Beek M, Blankenberg D, Bouvier D, Čech M, Chilton J, Clements D, Coraor N, Eberhard C, Grünig B, Guerler A, Hillman-Jackson J, Von Kuster G, Rasche E, Soranzo N, Turaga N, Taylor J, Nekrutenko A, Goecks J. 2016. The Galaxy platform for accessible, reproducible and collaborative biomedical analyses: 2016 update. *Nucleic Acids Res* 44:W3–W10. <https://doi.org/10.1093/nar/gkw343>.
- Goraichuk IV, Kulkarni AB, Williams-Coplin D, Suarez DL, Afonso CL. 2019. First complete genome sequence of currently circulating infectious bronchitis virus strain DMV/1639 of the GI-17 lineage. *Microbiol Resour Announc* 8:e00840-19. <https://doi.org/10.1128/MRA.00840-19>.
- Dimitrov KM, Sharma P, Volkening JD, Goraichuk IV, Wajid A, Rehmani SF, Basharat A, Shittu I, Joannis TM, Miller PJ, Afonso CL. 2017. A robust and cost-effective approach to sequence and analyze complete genomes of small RNA viruses. *Virology* 14:72. <https://doi.org/10.1186/s12985-017-0741-5>.
- Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* 25:1754–1760. <https://doi.org/10.1093/bioinformatics/btp324>.
- Ammayappan A, Upadhyay C, Gelb J, Vakharia VN. 2008. Complete genomic sequence analysis of infectious bronchitis virus Ark DPI strain and its evolution by recombination. *Virology* 5:157. <https://doi.org/10.1186/1743-422X-5-157>.
- Ammayappan A, Upadhyay C, Gelb J, Vakharia VN. 2009. Identification of sequence changes responsible for the attenuation of avian infectious bronchitis virus strain Arkansas DPI. *Arch Virol* 154:495–499. <https://doi.org/10.1007/s00705-009-0325-9>.