

Identification of Pantropic Canine Coronavirus in a Wolf (*Canis lupus italicus*) in Italy

Authors: Alfano, Flora, Dowgier, Giulia, Valentino, Maria Paola, Galiero, Giorgio, Tinelli, Antonella, et. al.

Source: Journal of Wildlife Diseases, 55(2) : 504-508

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/2018-07-182>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Identification of Pantropic Canine Coronavirus in a Wolf (*Canis lupus italicus*) in Italy

Flora Alfano,^{1,3} Giulia Dowgier,² Maria Paola Valentino,¹ Giorgio Galiero,¹ Antonella Tinelli,² Nicola Decaro,² and Giovanna Fusco¹ ¹Istituto Zooprofilattico Sperimentale del Mezzogiorno, Via della Salute 2, 80055 Portici (NA), Italy; ²Department of Veterinary Medicine, University of Bari, Valenzano (BA), Italy; ³Corresponding author (email: flora.alfano@cert.izsmportici.it)

ABSTRACT: We report a case in an Italian wolf (*Canis lupus italicus*) of pantropic canine coronavirus infection, which has previously been detected only in dogs. The wolf was coinfecting by canine parvovirus type 2b and canine adenovirus type 2, which highlighted the crucial role of epidemiologic surveys in European wild carnivores.

With few exceptions, populations of the European wolf (*Canis lupus*) have been increasing or stable during recent years, but their conservation is still threatened by illegal hunting, agricultural and silvicultural intensification, and land abandonment with reduction of free-ranging livestock (Votsi et al. 2016). Domestic dogs can transmit pathogens to wild carnivores through direct contact or indirect transmission through feces and urine (Millán et al. 2016). Several canine pathogens have been detected in European wolves including canine parvovirus (CPV), canine adenovirus types 1 (CAV-1) and 2 (CAV-2), and canine distemper virus (Martinello et al. 1997; Di Sabatino et al. 2014; Millán et al. 2016; Dowgier et al. 2018). Some of these pathogens were associated with high mortality and represented a severe threat to wolf conservation (Di Sabatino et al. 2014). Canine coronavirus (CCoV) causes mild enteritis in domestic dogs although a hypervirulent variant, named pantropic CCoV (pCCoV), is associated with mortality and leukopenia (Buonavoglia et al. 2006). To date, pCCoV infection has been reported only in dogs (Decaro and Buonavoglia 2011) and enteric CCoV has been only sporadically detected in wolves (Molnar et al. 2014). We report a case of pCCoV infection in an Italian wolf (*Canis*

lupus italicus) that was coinfecting with other canine viruses.

We submitted samples collected from various tissues (gut, heart, brain, spleen, liver, lungs) during the necropsy of a wolf found dead near Avellino (Italy), latitude 40°50'37 N, longitude 15°1'4 E, on December 2016, to RNA/DNA extraction by the QIASymphony DSP Virus/Pathogen kit (Qiagen, Hilden, Germany). Extracts were screened by gel-based or real-time PCR assays for canine viral pathogens including CPV, CCoV, CAV-1/CAV-2, canine distemper virus, canine alphaherpesvirus, and rotaviruses (Decaro et al. 2013). By molecular assays, the wolf was found to have a triple infection caused by CCoV, CPV-2b, and CAV-2. The CPV-2b strain was detected in several internal organs including gut, heart, brain, spleen, liver, and lungs whereas CAV-2 was recovered from lungs, spleen, and liver. The CCoV strain was detected in the gut and in nonintestinal tissues (heart, brain, spleen) and was typed as CCoV-IIa. According to previous studies (Decaro et al. 2013), the detection of CCoV-IIa in nonintestinal tissues allowed characterization of the strain as pCCoV.

The CCoV sequences were analyzed by the Geneious platform (version 10.1.3; Biomatters Ltd., Auckland, New Zealand) while similarity to GenBank sequences was assessed by BLAST (National Center for Biotechnology Information 2018) and FASTA (European Bioinformatics Institute 2018). The spike gene of the detected pCCoV strain, referred to as CCoV/wolf/2016/IT (GenBank accession no. MF991150), showed the highest degree of nucleotide (nt) identity (97%) with strain CCoV-IIa CCoV/dog/HCM27/2014 (GenBank no. LC190906) from a Vietnamese dog

(*Canis lupus familiaris*), followed by strains FCoV NTU156/P/2007 (GenBank no. GQ152141) from a Taiwanese cat (*Felis catus*) with 95% nt identity and GZ43/2003 (GenBank no. EF192155) from a Chinese raccoon dog (*Nyctereutes procyonoides*) with 93% nt identity. A lower degree of genetic relatedness was observed with CCoV-IIa strains from domestic dogs in China (86% nt identity): HLJ-071 (GenBank no. KY063616), HLJ-072 (no. KY063617), HLJ-073 (no. KY063618); Greece (86% nt identity): pCCoV 56/08 (no. GQ121370) and 59/08 (no. GQ121371); and Italy (85% nt identity): pCCoV CB/05 (no. KP981644).

Phylogenetic trees were generated from MAFFT Multiple Sequence Alignment Software (version 7; Biomatters Ltd.) by Geneious platform (version 10.1.3; using the neighbor joining and Bayesian methods). Bayesian inference was performed using four chains run over one million generations (with the first 2,000 trees discarded as burn-in) and supplying statistical support with subsampling over 200 replicates. According to the phylogenetic analysis (Figs. 1, 2), CCoV/wolf/2016/IT clusters with several CCoV-IIa and FCoV-II strains identified in European and extra-European countries, with no obvious geographic distribution. A pattern of node-based sub-clustering is displayed by both phylogenetic trees, showing that CCoV/wolf/2016/IT segregates with a Vietnamese strain (CCoV/dog/HCM27/2014) and a raccoon dog CCoV from China (GZ43/2003). Interestingly, other putative pCCoVs identified in previous studies were less-closely related to the wolf strain in comparison with enteric strains (Buonavoglia et al. 2006; Decaro et al. 2013).

Our findings showed that CCoV was circulating in the Italian wolf population, indicating that potentially fatal infections caused by pCCoV could be expected in this carnivore species. Interestingly, the wolf pCCoV strain was only distantly related to Italian CCoVs of canine origin, displaying higher genetic identities to strains circulating in Asian carnivores. It is therefore possible that the virus was introduced into Italy

through the importation of dogs or other carnivores from Asia, as suggested for other viral agents (Mira et al. 2018). Accordingly, CCoV infection is very common in domestic dogs, foxes, and raccoon dogs in China (Wang et al. 2006). While enteric CCoV causes self-limiting enteritis, pCCoV is responsible for fatal infections in dogs, indicating that this canine pathogen may represent an additional threat to the conservation of the Italian wolf population.

Two other canine viruses, CAHV-2 and CPV, were included in the screening due to a recent report of mortality in captive wolves (Dowgier et al. 2018). Millán et al. (2016) identified CAHV-2 in Iberian wolves (*Canis lupus signatus*) while CAHV-1 and CAHV-2 were recently detected in captive wolves (*Canis lupus lupus*) in France (Dowgier et al. 2018). To date, CAHV-1 (but not CAHV-2) has been detected in Italian wolves (Pizzurro et al. 2017), with the present study first reporting the detection of CAHV-2 in these wild carnivores.

Although CPV exposure of wolves had previously been reported in Italy (Martinello et al. 1997), our study provided molecular evidence for the active circulation of this virus. All the CPV strains isolated from wolves in Italy were identified as type 2b (Battilani et al. 2001), with the same virus strain circulating among dogs and wolves. Investigations on Iberian wolves (*Canis lupus signatus*) revealed that most of the wolf cases were due to a CPV-2c strain identical to that circulating in domestic dogs (Millán et al. 2016). This would support the idea that infection of wolves depends on periodic spillover from dogs. Indeed, previous studies suggested that dogs might be a reservoir for viral infections of wild canids (Ferreira et al. 2009; Molnar et al. 2014).

Given the possible negative impact of viral strains with increased pathogenicity on wolf populations, and because little information is available on pathogens infecting the Italian wolf, our results are important from the conservation perspective and highlight the need for continuous monitoring of pathogen circulation in wildlife.




FIGURE 1. Neighbor joining tree inferred from multiple nucleotide sequence alignment of the 5'-terminal region of the spike gene of the canine coronavirus (CCoV) strain detected in an Italian wolf (*Canis lupus italicus*) in 2016 and representative CCoVs, porcine transmissible gastroenteritis viruses (TGEVs), and feline coronaviruses (FCoVs) available on GenBank, recruited among different subgenotypes reported from different countries. For each sequence a short name, host species, collection date, and country are described in addition to




FIGURE 2. Bayesian phylogenetic tree inferred from multiple nucleotide sequence alignment of the 5'-terminal region of the spike gene of the canine coronavirus (CCoV) strain detected in an Italian wolf (*Canis lupus italicus*) in 2016 and representative CCoVs, porcine transmissible gastroenteritis viruses (TGEVs), and feline coronaviruses (FCoVs) available on GenBank, recruited among different subgenotypes reported from different countries. For each sequence a short name, host species, collection date, and country are described in addition to the GenBank accession numbers indicated in parentheses. Pantropic strains are defined as pCCoV whereas all the other CCoV are referred as type I or II based on the commonly accepted criteria. The scale bar indicates the estimated numbers of nucleotide substitutions per site. The distantly related *Betacoronavirus-1* canine respiratory coronavirus (CRCoV) 240/05 (DQ662401) was used as outgroup.

the GenBank accession numbers indicated in parentheses. Pantropic strains are defined as pCCoV whereas all the other CCoV are referred as type I or II based on the commonly accepted criteria. The scale bar indicates the estimated numbers of nucleotide substitutions per site. The distantly related *Betacoronavirus-1* canine respiratory coronavirus (CRCoV) 240/05 (DQ662401) was used as outgroup.

LITERATURE CITED

- Battilani M, Scagliarini A, Tisato E, Turilli C, Jacoboni I, Casadio R, Prosperi S. 2001. Analysis of canine parvovirus sequences from wolves and dogs isolated in Italy. *J Gen Virol* 82:1555–1560.
- Buonavoglia C, Decaro N, Martella V, Elia G, Campolo M, Desario C, Castagnaro M, Tempesta M. 2006. Canine coronavirus highly pathogenic for dogs. *Emerg Infect Dis* 12:492–494.
- Decaro N, Buonavoglia C. 2011. Canine coronavirus: Not only an enteric pathogen. *Vet Clin North Am Small Anim Pract* 41:1121–1132.
- Decaro N, Cordonnier N, Demeter Z, Egberink H, Elia G, Grellet A, Le Poder S, Mari V, Martella V, Ntasis V, et al. 2013. European surveillance for pantropic canine coronavirus. *J Clin Microbiol* 51:83–88.
- Di Sabatino D, Lorusso A, Di Francesco CE, Gentile L, Di Pirro V, Bellacicco AL, Giovannini A, Di Francesco G, Marruchella G, Marsilio F, et al. 2014. Arctic lineage-canine distemper virus as a cause of death in Apennine wolves (*Canis lupus*) in Italy. *PLoS One* 9:e82356.
- Dowgier G, Lahoreau J, Lanave G, Losurdo M, Varrello K, Lucente MS, Ventriglia G, Bozzetta E, Martella V, Buonavoglia C, et al. 2018. Sequential circulation of canine adenoviruses 1 and 2 in captive wild carnivores, France. *Vet Microbiol* 221:67–73.
- European Bioinformatics Institute. 2018. *FASTA protein similarity search*. <http://www.ebi.ac.uk/fasta33>. Accessed January 2018.
- Ferreyra H, Calderón MG, Marticorena D, Marull C, Leonardo BC. 2009. Canine distemper infection in crab-eating fox (*Cerdocyon thous*) from Argentina. *J Wildl Dis* 45:1158–1162.
- Martinello F, Galuppo F, Ostanello F, Guberti V, Prosperi S. 1997. Detection of canine parvovirus in wolves from Italy. *J Wildl Dis* 33:628–631.
- Millán J, López-Bao JV, García EJ, Oleaga Á, Llaneza L, Palacios V, de la Torre A, Rodríguez A, Dubovi EJ, Esperón F. 2016. Patterns of exposure of Iberian wolves (*Canis lupus*) to canine viruses in human-dominated landscapes. *Ecohealth* 13:123–134.
- Mira F, Purpari G, Lorusso E, Di Bella S, Gucciardi F, Desario C, Macaluso G, Decaro N, Guercio A. 2018. Introduction of Asian canine parvovirus in Europe through dog importation. *Transbound Emerg Dis* 65:16–21.
- Molnar B, Duchamp C, Möstl K, Diehl PA, Betschart B. 2014. Comparative survey of canine parvovirus, canine distemper virus and canine enteric coronavirus infection in free-ranging wolves of central Italy and south-eastern France. *Eur J Wildl Res* 60:613–624.
- National Center for Biotechnology Information. 2018. *Basic local alignment search tool (BLAST)*. <http://blast.ncbi.nlm.nih.gov/Blast.cgi>. Accessed January 2018.
- Pizzurro F, Marcacci M, Zaccaria G, Orsini M, Cito F, Rosamilia A, Di Renzo L, Malatesta D, Di Sabatino D, Lorusso A. 2017. Genome sequence of canine adenovirus type 1 isolated from a wolf (*Canis lupus*) in southern Italy. *Genome Announc* 5:e00225.
- Votsi NE, Zomeni MS, Pantis JD. 2016. Evaluating the effectiveness of Natura 2000 network for wolf conservation: A case-study in Greece. *Environ Manage* 57:257–270.
- Wang Y, Ma G, Lu C, Wen H. 2006. Detection of canine coronaviruses genotype I and II in raised Canidae animals in China. *Berl Munch Tierarztl Wochenschr* 119:35–39.

Submitted for publication 25 July 2018.

Accepted 14 August 2018.