

Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Journal Pre-proof

Oleoylethanolamide, A Bioactive Lipid Amide, as A Promising Treatment Strategy for Coronavirus/COVID-19

Samad Ghaffari, Neda Roshanravan, Helda Tutunchi, Alireza Ostadrahimi, Mahboub Pouraghaei, Behnam Kafil

PII: S0188-4409(20)30475-6

DOI: https://doi.org/10.1016/j.arcmed.2020.04.006

Reference: ARCMED 2491

To appear in: Archives of Medical Research

Received Date: 5 April 2020

Revised Date: 8 April 2020

Accepted Date: 10 April 2020

Please cite this article as: Ghaffari S, Roshanravan N, Tutunchi H, Ostadrahimi A, Pouraghaei M, Kafil B, Oleoylethanolamide, A Bioactive Lipid Amide, as A Promising Treatment Strategy for Coronavirus/ COVID-19, *Archives of Medical Research* (2020), doi: https://doi.org/10.1016/j.arcmed.2020.04.006.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Copyright © 2020 Published by Elsevier Inc. on behalf of IMSS.



E20_423

Archives of Medical Research 51 (2020) x-x

OPINION

Oleoylethanolamide, A Bioactive Lipid Amide, as A Promising Treatment Strategy for Coronavirus/COVID-19

Samad Ghaffari,^a Neda Roshanravan,^a Helda Tutunchi,^b Alireza Ostadrahimi,^b Mahboub Pouraghaei,^c and Behnam Kafil^b

^aCardiovascular Research Center, Tabriz University of Medical Sciences, Tabriz, Iran ^bNutrition Research Center, Department of Clinical Nutrition, School of Nutrition and Food Sciences, Tabriz University of Medical Sciences, Tabriz, Iran ^cEmergency Medicine Research Team, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

Received for publication April 5, 2020; accepted April 9, 2020 (ARCMED_2020_423).

Address reprint requests to: Neda Roshanravan, Dr., Cardiovascular Research Center, Tabriz University of Medical Sciences, Pstal Code 5166615573, Tabriz, Iran; Phone and Fax: (0098) (41) 33373919; E-mail: Neda.roshanravan10@gmail.com

The current outbreak of COVID-19 (coronavirus) has been identified by World Health Organization

E20_423

(WHO) as a global pandemic. With the emergence of the COVID-19 virus and considering the lack of effective pharmaceutical treatment for it, there is an urgent need to identify safe and effective drugs or potential adjuvant therapy in this regard. Bioactive lipids with an array of known healthpromoting properties can be suggested as effective agents in alleviating acute respiratory stress induced by virus. The bioactive lipid amide, oleoylethanolamide (OEA), due to several distinctive homeostatic properties, including anti-inflammatory activities, modulation of immune response, and anti-oxidant effects can be considered as a novel potential pharmacological alternative for the management of COVID-19.

Key Words: Bioactive lipids, Coronavirus, COVID-19, OEA, Oleoylethanolamide.

Introduction

SARS-CoV-2: Prevalence, Phylogenetics, Taxonomy and Structural Biology

The novel coronavirus (COVID-19) has led to the economic disruptions and global health concerns due to its sustained human-to-human transmission and rapid spread (1). As of 1 April 2020, this severe acute respiratory syndrome (SARS-CoV-2) has affected a total of 867,922 confirmed cases with 43,152 deaths in 200 countries and territories (2). The acute respiratory distress syndrome (ARDS), which results in multiple organ failure and sepsis, remains the most common cause of death in these patients. Older adults, subjects with underlying chronic illnesses including cancer, hypertension and diabetes are the most seriously affected groups (3). There is currently no known treatment for SARS-CoV-2 infection, and investigators in many countries around the world have

E20_423

undertaken several clinical trials to solve the problem.

This highly transmittable and pathogenic viral belongs to the subfamily Orthocoronavirinae in the family of Coronaviridae, in the order Nidovirales (4). The subfamily of coronaviruses family includes alpha (α), beta (β), gamma (γ) and delta (δ) coronavirus (5). Coronaviruses were initially thought to infect only animals, but evidences from the last decade indicate that the disease is capable of infecting humans as well. SARS outbreak caused by SARS-CoV in 2002 in Guangdong, China, and Middle East Respiratory Syndrome Coronavirus (MERS-CoV) prevalence in 2012 in Saudi Arabia, were approval for the possibility of the virus spread in humans and the witness for the lethality of coronaviruses (6,7).

Genetic recombination event at S protein in the receptor binding domain (RBD) region of SARS-CoV-2 is thought to be the most important reason for the higher transmission rate of SARS-CoV-2 regarding to SARS-CoV. The SARS-CoV-2, a large sized virus (approximately 120 nanometers in diameter), is a positive-sense single-stranded RNA (+ssRNA) virus, and contains an outer lipid (fatty) membrane. Its RNA sequence is approximately 30,000 bases in length (8). An analysis of genome sequence data from SARS-CoV-2 revealed that SARS-CoV-2 was approximately 79% similar to SARS-CoV at the nucleotide sequence. Interestingly, new evidence suggests that the SARS-CoV-2 RBD is well suited for binding to the human angiotensin-converting enzyme 2 (ACE2) receptors, which are expressed in the membranes of various cells in the body, such as type II alveolar epithelial cells of the lungs. In fact, binding of the SARS-CoV to the ACE2 (9) receptors, disrupts the function of it (10). Intestine, kidney and blood vessels are the other main cells for ACE2 expression, and this fact may explain why some patients with Covid-19 experience gastrointestinal symptoms (11). The virus binds to the host-cell ACE2 receptor via a particular surface glycoprotein called a "spike" (peplomer), and enters the host cell. It is supposed that the entry of the virus to human cells increases

E20_423

the inflammatory activity through various mechanisms with consequent release of pro-inflammatory cytokines, which cause serious damage, specially to the respiratory tract (12). Theoretically, it is assumed that a decrease in the activity of ACE2 in cell membranes may reduce the ability of SARS-CoV-2 to penetrate cells (13).

Along with other endemic human coronaviruses, it seems that the SARS-CoV-2 will become the fifth endemic coronavirus in the human population. Nowadays, discovering therapeutic options from currently available agents appear to be essential for the treatment and prophylaxis of this pandemic. In this manuscript, we aimed to introduce oleoylethanolamide (OEA), a bioactive lipid mediator, as a novel potential pharmacological alternative for the management of COVID-19.

Oleoylethanolamide

The bioactive lipid amide OEA is synthesized in the gastro-intestinal tract, and is related to several distinctive homeostatic properties, including anti-inflammatory activities, immune response, stimulation of lipolysis and fatty acid oxidation (14). OEA, a member of the N-acylethanolamine (NAE) family, is derived from the omega-9 monounsaturated fatty acid, oleic acid. Previous studies have indicated that the down-regulation of OEA levels arises in situation such as exposure to stress, which contributes to the increase in inflammatory markers and the NAE catabolism (15,16). In the current epidemiological studies on inflammatory related diseases, OEA is considered an endocannabinoid like lipid, which interacts with the peroxisome proliferator-activated receptor- α (PPAR- α) and mediates the anti-inflammatory processes (17). It is generally accepted that the endocannabinoid system (ECS) consists of the membrane cannabinoid receptors (cannabinoid receptor type 1 [CB1R] and type 2 [CB2R]), endogenous ligands (endcannabinoids), and enzymes

E20_423

responsible for the synthesis and degradation of ligands (18). The collaboration of the ECS in the management and elimination of infectious agents such as viruses, bacteria, and some protozoa is indicated previously (19). Scientific researchers recently showed that the activation of the ECS relieves pain and reduces inflammation in the lungs. Based on their evidence, endogenous and exogenous cannabinoids can be considered as therapeutic potential agents for respiratory pathogen clearance (20).

Oleoylethanolamide and SARS-CoV-2 Infection

Based on previous studies, angiotensin receptors can be inhibited by several unsaturated fatty acids and their metabolites such as oleic acid. Emerging evidence suggests that these fatty acids can decrease the affinity of angiotensin receptors (21). Recent evidences also indicate that some unsaturated fatty acids can be served as endogenous anti-viral compounds, and their deficiency make humans more susceptible to certain viral infections including SARS-CoV-2, SARS and MERS (22). In view of the fact that SARS-CoV-2 infection leads to increased release of the pro-inflammatory cytokines, including interleukin-6 (IL-6) and IL-1 β in COVID-19 patients via binding to the Toll Like Receptors (TLRs) (12), it is assumed that OEA inhibits this pathway through its antiinflammatory properties. In fact, OEA binds with high affinity to PPAR- α receptors and initiates a cascade of events, which can eventually attenuate the inflammatory responses.

TLRs, important components of the innate immune system, are distinct classes of pattern-recognition receptors (PRRs), and are localized in the cell surface or in the intracellular compartments such as endolysosome, endoplasmic reticulum, endosome, and lysosomes. They can detect viral danger signals in the extracellular milieu and endosomes by leucine rich repeat-containing trans membrane

E20_423

proteins in their structure (23). The initiation of this downstream signaling, culminates in the activation of transcription nuclear factor kappa B (NF- κ B), and leads to the over expression of inflammatory cytokines adhesion molecules and chemokines (24,25).

Peroxisome proliferator-activated receptors (PPARs) are other classes of transcriptional factors involved in modulation of inflammatory pathways. These nuclear receptor proteins consist of three subtypes including PPAR- α , PPAR- γ , and PPAR- β/δ (25). PPARs can alleviate the expression of inflammatory mediators via inducing IkB α as the main inhibitor of NF- κ B signaling pathway (26). Previously, it has been indicated that the activation of TLRs pathways can attenuate the expression level of PPARs, and intensify the expression level of pro-inflammatory cytokines such as tumor nex crosis factor- α (TNF- α) (27).

Interestingly, OEA may modulate cross-talk between PPAR- α and TLRs and regulates the inflammatory responses in COVID-19. In other words, OEA may have a beneficial synergistic effect against SARS-CoV-2 infection. OEA, a high affinity endogenous ligand of PPAR- α (28), binds to PPAR- α receptors, and increases the expression level of anti-inflammatory cytokine such as IL-10. In addition, it attenuates the inflammatory responses and decreases the expression of TLR4, and interfering with the ERK1/2/AP-1/STAT3 signaling cascade (29–31). In a recent clinical trial, OEA supplementation could decrease inflammation in obese patients via reducing serum concentrations of inflammatory markers including IL-6 and TNF- α (32). On the whole, the results found in previous studies demonstrate that the activation of PPAR- α by OEA can prevent the expression of genes coding for several inflammatory cytokines. It is well established that OEA can also significantly reduce lipopolysaccharide-induced oxidative/nitrosative stress and prevents endothelial cell damage. By increasing the activity of the anti-oxidative enzymes, these beneficial effects of OEA can be justified (33,34). It has been suggested that excessive inflammation, oxidation and an increased

E20_423

immune response are the main contributors to COVID-19 pathology. OEA through the antioxidant and anti-inflammatory effects may be effective in the attenuation of inflammation and oxidation in coronavirus infected subjects.

OEA and Safety

The utmost concern about OEA supplementation in COVID-19 patients is its safety. In previous clinical trials, doses of 250 mg/d of OEA oral intake by participants exhibited satisfactory safety when compared to placebo. Also, even when OEA was given to humans at a dose of 250 mg/d for 3 months, no side effects were reported (28,32).

Conclusion

Based on available evidences, exogenous administration of OEA appears to be a homeostatic signal to counter COVID-19 infection and alleviate patients' inflammatory status. At present, we have decided to conduct a clinical trial on COVID-19 patients in Iran to evaluate this hypothesis. We hope that the results of this trial provide new insight for researchers to tackle the virus.

Conflict of Interest

No potential conflict of interest disclosed.

Funding

E20_423

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Acknowledgements

The authors would like to thank the Cardiovascular Research Center of Tabriz University of Medical Sciences.

Reference

1. Shen K, Yang Y, Wang T, et al. Diagnosis, treatment, and prevention of 2019 novel coronavirus infection in children: experts' consensus statement. World J Pedtatr 2020. doi: 10.1007/s12519-020-00343-7. [Epub ahead of print].

 Organization WH. Coronavirus disease 2019 (COVID-19): situation report, 59. 2020 (Accessed March 20, 2020).

3. Wang A, Zhao W, Xu Z, et al. Timely blood glucose management for the outbreak of 2019 novel coronavirus disease (COVID-19) is urgently needed. Diabetes Res Clin Pract 2020;162:108118.

4. Shereen MA, Khan S, Kazmi A, et al. COVID-19 infection: origin, transmission, and characteristics of human coronaviruses. J Adv Res 2020;24:91–98.

E20_423

5. Banerjee A, Kulcsar K, Misra V, et al. Bats and coronaviruses. Viruses 2019;11:41.

6. Zhong N, Zheng B, Li Y, et al. Epidemiology and cause of severe acute respiratory syndrome (SARS) in Guangdong, People's Republic of China, in February, 2003. Lancet 2003;362:1353–1358.

 Schoeman D, Fielding BC. Coronavirus envelope protein: current knowledge. Virol J 2019;16:69.

Zhu N, Zhang D, Wang W, et al. A novel coronavirus from patients with pneumonia in China,
2019. N Engl J Med 2020;382:727–733.

 Zhang Y-Z, Holmes EC. A Genomic Perspective on the Origin and Emergence of SARS-CoV-2. Cell 2020;181:1–5.

 Diaz JH. Hypothesis: angiotensin-converting enzyme inhibitors and angiotensin receptor blockers may increase the risk of severe COVID-19. J Travel Med. 2020. pii: taaa041. doi: 10.1093/jtm/taaa041. [Epub ahead of print].

11. Monteleone G, Ardizzone S. Are patients with inflammatory bowel disease at increased risk for Covid-19 infection? J Crohns Colitis 2020. pii: jjaa061. doi: 10.1093/ecco-jcc/jjaa061. [Epub ahead of print].

E20_423

12. Conti P, Ronconi G, Caraffa A, et al. Induction of pro-inflammatory cytokines (IL-1 and IL-6) and lung inflammation by COVID-19: anti-inflammatory strategies. J Biol Regul Homeost Agents. 2020;34(2). pii: 1. doi: 10.23812/CONTI-E. [Epub ahead of print].

 Letko M, Marzi A, Munster V. Functional assessment of cell entry and receptor usage for SARS-CoV-2 and other lineage B betacoronaviruses. Nat Microbiol 2020;5:562–569.

14. Di Paola M, Bonechi E, Provensi G, et al. Oleoylethanolamide treatment affects gut microbiota composition and the expression of intestinal cytokines in Peyer's patches of mice. Sci Rep 2018;8:1–12.

15. Bowen KJ, Kris-Etherton PM, Shearer GC, et al. Oleic acid-derived oleoylethanolamide: a nutritional science perspective. Prog Lipid Res 2017;67:1–15.

 Tutunchi H, Asghari Jafarabadi M, Hoojeghani S, et al. General and abdominal obesity is related to socioeconomic status and food choices: a cross-sectional study. Nutrition and Food Science.
2019;50:61–73.

17. Tutunchi H, Saghafi-Asl M, Ostadrahimi A. A systematic review of the effects of oleoylethanolamide, a high affinity endogenous ligand of PPAR-alpha, on the management and prevention of obesity. Clin Exp Pharmacol Physiol 2020;47:543–552.

18. Chye Y, Christensen E, Solowij N, et al. The endocannabinoid system and cannabidiol's promise

E20_423

for the treatment of substance use disorder. Front Psychiatry 2019;10:63.

19. Hernández-Cervantes R, Méndez-Díaz M, Prospéro-García Ó, et al. Immunoregulatory role of cannabinoids during infectious disease. Neuroimmunomodulation 2017;24:183–199.

20. Turcotte C, Blanchet M-R, Laviolette M, et al. Impact of cannabis, cannabinoids, and endocannabinoids in the lungs. Front Pharmacol 2016;7:317.

21. Goodfriend TL, Ball DL. Fatty acid effects on angiotensin receptors. J Cardiovasc Pharmacol 1986;8:1276–1283.

22. Das UN. Can Bioactive Lipids Inactivate Coronavirus (COVID-19)? Arch Med Res 2020.

23. Akira S, Uematsu S, Takeuchi O. Pathogen recognition and innate immunity. Cell.2006;124:783–801.

24. Barton GM, Medzhitov R. Toll-like receptor signaling pathways. Science 2003;300:1524–1525.

25. Lee SM, Kok K-H, Jaume M, et al. Toll-like receptor 10 is involved in induction of innate immune responses to influenza virus infection. Proc Natl Acad Sci 2014;111:3793–3798.

26. Youssef J, Badr M. Role of peroxisome proliferator-activated receptors in inflammation control.Biomed Res Int 2004;2004:156–166.

E20_423

27. Necela BM, Su W, Thompson EA. Toll \Box like receptor 4 mediates cross \Box talk between peroxisome proliferator \Box activated receptor γ and nuclear factor \Box κ B in macrophages. Immunology 2008;125:344–358.

28. Tutunchi H, Ostadrahimi A, Saghafi-Asl M, et al. Oleoylethanolamide supplementation in obese patients newly diagnosed with non-alcoholic fatty liver disease: effects on metabolic parameters, anthropometric indices, and expression of PPAR- α , UCP1, and UCP2 genes. Pharmacol Res 2020;156:104770.

29. Tutunchi H, Ostadrahimi A, Saghafi \Box Asl M, et al. The effects of oleoylethanolamide, an endogenous PPAR $\Box \alpha$ agonist, on risk factors for NAFLD: A systematic review. Obes Rev 2019;20:1057–1069.

30. Yang L, Guo H, Li Y, et al. Oleoylethanolamide exerts anti-inflammatory effects on LPSinduced THP-1 cells by enhancing PPAR α signaling and inhibiting the NF- κ B and ERK1/2/AP-1/STAT3 pathways. Sci Rep 2016;6:34611.

31. Antón M, Rodríguez González A, Rodríguez Rojo IC, et al. Increased plasma oleoylethanolamide and palmitoleoylethanolamide levels correlate with inflammatory changes in alcohol binge drinkers: the case of HMGB1 in women. Addict Biol 2018;23:1242–1250.

32. Payahoo L, Khajebishak Y, Jafarabadi MA, et al. Oleoylethanolamide supplementation reduces

E20_423

inflammation and oxidative stress in obese people: a clinical trial. Adv Pharm Bull 2018;8:479.

33. Ma L, Guo X, Chen W. Inhibitory effects of oleoylethanolamide (OEA) on H2O2-induced human umbilical vein endothelial cell (HUVEC) injury and apolipoprotein E knockout (ApoE–/–) atherosclerotic mice. Int J Clin Exp Pathol 2015;8:6301.

34. Zhang R, Wang X, Ni L, et al. COVID-19: Melatonin as a potential adjuvant treatment. Life Sci. 2020:117583.

Rect

E20_423

NAME FOR TOP OF PAGES: Ghaffari, et al./Archives of Medical Research 51 (2020) x-x

Running title: Oleoylethanolamide and Coronavirus/COVID-19

Journal Prevention

- Outbreak of COVID-19 (coronavirus) has been identified by World Health Organization as a global pandemic.
- There is an urgent need to identify safe and effective drugs or potential adjuvant therapy in this regard.
- Oleoylethanolamide (OEA) may be considered as a novel potential adjuvant pharmacological alternative for the management of COVID-19.
- OEA via its antioxidant and anti-inflammatory properties may be effective in the management of coronavirus infected subjects.

Journal Prever