




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Contents lists available at ScienceDirect

Oral Oncology

journal homepage: www.elsevier.com/locate/oraloncology

Letter to the editor

A probable surge in oral lichen planus cases under the aura of coronavirus in females in India

A mucocutaneous disorder, Oral Lichen Planus (OLP) is seen to affect the oral cavity commonly. As it does not fit into the attire of a classical autoimmune disorder, its best portrayed as a chronic inflammatory disease. It is mediated immunologically by the action of cytotoxic T-cells directed against basal keratinocytes causing epithelial damage [1]. The commonest part of oral cavity involved is the gingiva and tongue with a high predilection for females in between fourth and fifth decade of age. LP is found to be significantly associated with an increased risk of cancer of lip, tongue, oralcavity [2], OLP, in itself is a very painful disease with severe burning symptoms in the oral cavity for the affected person. After taking medication for long time too, it recurs most of the time. This again induces fear amongst the affected of developing an incurable disease. Stress has been infamously branded to alter several parameters of the endocrine system and immune response in individuals with OLP. Even higher rate of sleep problems, depression and anxiety has been stated in OLP patients [3]. A connotation between psychological disorders and the development of OLP has already been established through studies [4].

Majority of Indian woman, are exposed to various cultural and moral attributes at an early age. Though modernisation in a developing country like India is revealing them as epitome of courage, resilience and the ability to fit into diverse roles, still anxiety and stress grips them more firmly [5]. Understanding stress pathogenesis, it can activate the sympathetic nervous system and the hypothalamic pituitary adrenocortical axis. This leads to heavy influx of inflammatory cytokines like IL-6, IL-1 and TNF by the sympathetic nervous system. Modifications in the kynurenine metabolites as a result of this influx leads to neurotoxic changes in brain. The excess secretion of cortisol originates as immunosuppression through multiple mechanisms (NF κ B, AP1, Raf, MAPK mediated signalling pathways). Even, hormonal changes in females makes them prone to stress induced systemic disorders. Studies have also shown females to be more susceptible to the central effects of increased cytokines during stress [6].

The World Health Organization (WHO) on March 11, 2020, declared the novel coronavirus (COVID-19) outbreak a global pandemic by WHO Director-General, Dr. Tedros Adhanom Ghebreyesus. Today, India is under a 21-day nationwide lockdown starting from 25th March in a desperate attempt to contain the spread of novel coronavirus SARS-Cov-2, which has already broken the 7000-mark in the country [7]. As people are being placed in quarantine, isolation and travel restrictions, surge in fear and chaos is evident among the mass. The psychological toll is complex related to factors like fear of losing loved ones, fear of losing livelihood, compulsive need to hoard, phobia for contracting the infection, etc. The disastrous effect of coronavirus on human morbidity and mortality is evident from reports from other countries. An impression of triggering potential mayhem for a low socio-economic country like India is in abundance. The lock down has literally bought down the Indian economy to a standstill having huge im-

pact on Indian population. Its leading to superior levels of anxiety and sense of insecurity amongst the people, especially the lower socioeconomic classes. This, pandemic in itself is a colossal causative factor for stress and anxiety amongst Indian women. Though the malignant transformation from OLP was assessed to be 0.5% in a recent systematic review and meta-analysis [9], but the question remains; What are the chances of a surge in oral lichen planus cases under the impact of this pandemic stress on females in India? Will this stress act an inducing or as a contributing factor for the transformation, if at all? The ICD-11 has recognized a new group of 'disorders specifically associated with stress' that identifies disorders in which external stress is a necessary and prominent causal factor [8]. Prioritising public health, especially in context to mental health is the need of the hour and the key to wellbeing in this era of pandemic [10].

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

- [1] Hirschfeld J, Higham J, Chatzistavrianou D, Blair F, Richards A, Chapple ILC. Systemic disease or periodontal disease? Distinguishing causes of gingival inflammation: a guide for dental practitioners. Part 1: immune-mediated, autoimmune-inflammatory, and hereditary lesions. *Br Dent J* 2019;227(11):961–6.
- [2] Halonen P, Jakobsson M, Heikinheimo O, Riska A, Gissler M, Pukkala E. Cancer risk of Lichen planus: A cohort study of 13,100 women in Finland. *Int J Cancer* 2018;142(1):18–22.
- [3] Adamo D, Ruoppo E, Leuci S, Aria M, Amato M, Mignogna MD. Sleep disturbances, anxiety and depression in patients with oral lichen planus: a case-control study. *J Eur Acad Dermatol Venereol* 2015;29(2):291–7.
- [4] Cerqueira JDM, Moura JR, Arsati F, Lima-Arsati YBDO, Bittencourt RA, Freitas VS. Psychological disorders and oral lichen planus: A systematic review. *J Invest Clin Dent* 2018;9:e12363.
- [5] De Sousa D, De Sousa A. Indian Womanhood: Some Psychological Concepts. *Mens Sana Monogr* 2015;13(1):16–22.
- [6] Albert PR. Why is depression more prevalent in women. *J Psychiatry Neurosci* 2015;40(4):219–21.
- [7] Banerjee D. How COVID-19 is overwhelming our mental health doi: 10.1038/nindia.2020.46.
- [8] Stein DJ, Szatmari P, Gaebel W, Berk M, Vieta E, Maj M, et al. Mental, behavioral and neurodevelopmental disorders in the ICD-11: an international perspective on key changes and controversies. *BMC Med* 2020;18:21.
- [9] Idrees M, Kujan O, Shearston K, Farah CS. Oral lichen planus has a very low malignant transformation rate: A systematic review and meta-analysis using strict diagnostic and inclusion criteria. *J Oral Pathol Med* 2020. <https://doi.org/10.1111/jop.12996>.
- [10] Send in the therapists? *The Lancet Psychiatry* 2020; 7(4): P291.

Samapika Routray ((MDS; Ph.D))* , Priyadarshini Mishra ((MD))
All India Institute of Medical Sciences, Bhubaneswar, India
E-mail address: drroutray.samapika@gmail.com (S. Routray).

* Corresponding author at: Department of Dentistry, All India Institute of Medical Sciences, Bhubaneswar 751019, Odisha, India.