




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Journal Pre-proof

COVID-19 pandemic and the stethoscope: don't forget to sanitize



Mark A. Marinella MD, FACP

PII: S0147-9563(20)30112-6
DOI: <https://doi.org/10.1016/j.hrtlng.2020.03.017>
Reference: YMHL 1619

To appear in: *Heart & Lung*

Received date: 27 March 2020
Accepted date: 30 March 2020

Please cite this article as: Mark A. Marinella MD, FACP , COVID-19 pandemic and the stethoscope: don't forget to sanitize, *Heart & Lung* (2020), doi: <https://doi.org/10.1016/j.hrtlng.2020.03.017>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier Inc.

COVID-19 pandemic and the stethoscope: don't forget to sanitize

Mark A. Marinella, MD, FACP

Associate Clinical Professor of Medicine, Wright State University School of Medicine, Dayton, OH

CORRESPONDENCE: Mark A. Marinella MD FACP, Miami Valley Hospital South, Comprehensive Cancer Center, 1500 Miami Valley Drive, Dayton, OH 45459, PHONE: 937 542 1407, FAX: 937 832 5344, MMarinella@daytonphysicians.com, 937-542-1407

The SARS-CoV-2 (COVID 19) pandemic has resulted in unprecedented rapid generation and dissemination of published data in short-order, in almost “real-time.” Thanks to rapid review and publication, as a medical community we are rapidly learning more of the epidemiology, clinical manifestations, complications, and potential reservoirs of SARS-CoV-2 that may enhance transmission. Clinicians in all specialties are able to keep current due to this rapid turn-around of essential information. Providers who work on the “front-lines” as first-responders or in the emergency department, as well as in hospital wards, the intensive care unit, or the outpatient setting, need to implement a variety of mechanisms to help curb the spread of this viral pandemic.

Indirect pathogen transmission from inanimate objects is of potential concern not only for the general public, but also for healthcare professionals whose hands come into frequent contact with hard surfaces. Indeed, nosocomial infectious outbreaks from bacterial contamination of thermometers¹ and blood pressure cuffs² have been reported. Viral pathogens have also been isolated on hard surfaces in the healthcare setting,^{3,4} but have not

received as much attention as a risk for nosocomial and person-to-person transmission until very recently with the COVID-19 pandemic.

In addition to efficient spread *via* respiratory secretions/droplets, SARS-CoV-2 can survive on solid surfaces found in the healthcare setting.⁵ van Doremalen *et al.* elegantly demonstrated that SARS-CoV-2 can remain viable, and thus potentially transmissible, for prolonged periods on stainless steel and plastic, and to a lesser extent, cardboard.⁵ This has raised concern regarding nosocomial transmission and supports public health officials' urging of the public to sanitize their hands frequently when coming into contact with public surfaces. Other coronaviruses such as the Middle East Respiratory Syndrome (MERS) coronavirus and SARS-CoV-1, have been reported to remain viable on inanimate surfaces such as glass, plastic or metal for up to nine days, but can be inactivated by alcohol solutions, hydrogen peroxide, and sodium hypochlorite.^{6,7}

Coronavirus viability could, in turn, result in self-innoculation if one touches their face or transmission to another person *via* hand contact.⁷

Healthcare personnel constantly come into contact with countertops, computers, phones, doorknobs, hospital charts, and stethoscopes, which have been shown to be colonized with a variety of bacterial pathogens.⁸ The stethoscope is utilized by countless healthcare personnel in various settings and has been shown to harbor potentially pathogenic bacteria and transmit bacteria to human skin.⁹ Cleaning the plastic diaphragm with standard cleaning solutions found in the healthcare setting can decrease bacterial load.⁹ Respiratory syncytial virus (RSV) has been shown to survive on stethoscope diaphragms and eradicated with

standard alcohol wipes.⁴ The severe fever with thrombocytopenia syndrome (SFTS) virus, an emerging fatal viral hemorrhagic fever in East Asia, has been recovered from stethoscopes and other hard surfaces in patient rooms who were diagnosed with SFTS, also raising concern for nosocomial transmission of highly pathogenic viruses.³ In light of the recent data that SARS-CoV-2 can survive on plastic surfaces,⁵ it is important for healthcare workers to clean the plastic stethoscope diaphragm at the same time as sanitizing their hands. Furthermore, “community” stethoscopes used by multiple individuals and sharing of stethoscopes should be avoided, if possible, in the midst of this pandemic in order to prevent nosocomial spread and potential self-inoculation. Although the stethoscope is a small piece of our equipment, as healthcare professionals, we can employ small maneuvers in attempt to curtail the spread of this virus, and this includes regular sanitizing of our stethoscopes.

References

1. Ilvornese LL, Dias S, Samel C, et al. Hospital-acquired infection with vancomycin-resistant *Enterococcus faecium* transmitted by electronic thermometers. *Ann Intern Med* 1992;117:112-16
2. Layton MC, Perez M, Heald P, et al. An outbreak of mupirocin-resistant *Staphylococcus aureus* on a dermatology ward with an environmental reservoir. *Infect Control Hosp Epidemiol* 1993;14:369-375
3. Ryu BH, Kim JY, Kim T, et al. Extensive severe fever with thrombocytopenia syndrome virus contamination in surrounding environment in patient rooms. *Clin Microbiol Infect* 2018;24:911.e1-911.e4

4. Blydt-Hanson T, Kanta S, Quennec PRT, et al. Recovery of respiratory syncytial virus from stethoscopes by conventional viral culture and polymerase chain reaction. *Pediatr Infect Dis J* 1999;18:164-5
5. Van Doremalen N, Bushmaker T, Morris DH, et al. Aerosol and surface stability of SARS-CoV-2 as compared with SARS-CoV-1. *N Engl J Med* DOI:10.1056/NEJMc2004973
6. Kampf G, Todt D, Pfaender S, et al. Persistence of coronaviruses on inanimate surfaces and their inactivation with biocidal agents. *J Hosp Infect* 2020;104:246-251
7. Otter JA, Donskey C, Yezli S, et al. Transmission of SARS and MERS coronaviruses and influenza virus in healthcare settings: the possible role of dry surface contamination. *J Hosp Infect* 2016;92:235-50
8. Marinella MA, Elder BL. Bacterial contamination of hospital charts. *Infect Dis Clin Pract* 2000;9:39-40
9. Marinella MA, Pierson C, Chenoweth C. The stethoscope: a potential source of nosocomial infection? *Arch Intern Med* 1997;157:786-90