



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

# Journal Pre-proof

Reorganization of Medical Oncology Departments during COVID-19 Pandemic: a Nationwide Italian Survey

Alice Indini, MD, Carlo Aschele, MD, PhD, Daniele Bruno, MD, Luigi Cavanna, MD, Mario Clerico, MD, Giammaria Fiorentini, MD, Luisa Fioretto, MD, Monica Giordano, MD, Vincenzo Montesarchio, MD, Cinzia Ortega, MD, Graziella Pinotti, MD, Alberto Scanni, MD, Claudio Zamagni, MD, Livio Blasi, MD, Francesco Grossi, MD



PII: S0959-8049(20)30165-9

DOI: <https://doi.org/10.1016/j.ejca.2020.03.024>

Reference: EJC 11384

To appear in: *European Journal of Cancer*

Received Date: 31 March 2020

Revised Date: 31 March 2020

Accepted Date: 31 March 2020

Please cite this article as: Indini A, Aschele C, Bruno D, Cavanna L, Clerico M, Fiorentini G, Fioretto L, Giordano M, Montesarchio V, Ortega C, Pinotti G, Scanni A, Zamagni C, Blasi L, Grossi F, Reorganization of Medical Oncology Departments during COVID-19 Pandemic: a Nationwide Italian Survey, *European Journal of Cancer*, <https://doi.org/10.1016/j.ejca.2020.03.024>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Elsevier Ltd. All rights reserved.

## **Reorganization of Medical Oncology Departments during COVID-19 Pandemic: a Nationwide Italian Survey.**

Alice Indini<sup>1</sup>, MD; Carlo Aschele<sup>2</sup>, MD, PhD; Daniele Bruno<sup>3</sup>, MD; Luigi Cavanna<sup>4</sup>, MD; Mario Clerico<sup>5</sup>, MD; Giammaria Fiorentini<sup>6</sup>, MD; Luisa Fioretto<sup>7</sup>, MD; Monica Giordano<sup>8</sup>, MD; Vincenzo Montesarchio<sup>9</sup>, MD; Cinzia Ortega<sup>10</sup>, MD; Graziella Pinotti<sup>11</sup>, MD; Alberto Scanni<sup>12</sup>, MD; Claudio Zamagni<sup>13</sup>, MD; Livio Blasi<sup>14</sup>, MD; Francesco Grossi<sup>1</sup>, MD.

<sup>1</sup>Medical Oncology Unit, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milan, Italy.

<sup>2</sup>Medical Oncology Department, Ospedale Sant'Andrea, La Spezia, Italy; Regional Coordinator of CIPOMO Liguria; Member of the CIPOMO National Directive Board.

<sup>3</sup>Oncology Unit, Ospedale del mare, Naples, Italy; Member of the CIPOMO National Directive Board.

<sup>4</sup>Oncology-Hematology Department, Hospital of Piacenza, Piacenza, Italy; Member of the CIPOMO National Directive Board.

<sup>5</sup>Medical Oncology Department ASL Biella, Biella, Italy; Past President of CIPOMO.

<sup>6</sup>Department of Oncology and Hematology, Ospedali Riuniti Marche Nord, Pesaro, Italy; Auditor of CIPOMO.

<sup>7</sup>Department of Oncology, SM Annunziata Hospital, Florence, Italy; Treasurer of CIPOMO.

<sup>8</sup>Medical Oncology Division, ASST-Lariana, Como, Italy; Regional Coordinator of CIPOMO Lombardia; Member of the CIPOMO National Directive Board.

<sup>9</sup>Oncology Unit, "Monaldi" Hospital, Naples, Italy; Chairman of CIPOMO.

<sup>10</sup>Department of Medical Oncology, Ospedale S. Lazzaro ASL CN2 Alba-Bra, Cuneo, Italy; Auditor of CIPOMO.

<sup>11</sup>Medical Oncology Department, ASST Sette Laghi, Varese, Italy; Vice President of CIPOMO.

<sup>12</sup>Department of Medical Oncology, Fatebenefratelli e Oftalmico Hospital, Milan, Italy; President Emeritus of CIPOMO.

<sup>13</sup>Medical Oncology Unit, S.Orsola-Malpighi Hospital, Bologna, Italy; Member of the CIPOMO National Directive Board.

<sup>14</sup> Medical Oncology Unit, ARNAS Civico, Palermo, Italy; President of CIPOMO.

### **Corresponding author:**

Dr. Alice Indini,

Medical Oncology Unit, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico,

Via Francesco Sforza 35, 20122, Milano, Italy

Phone: +39-0255035558; fax: +39-0255032659

E-mail: alice.indini@gmail.com

**Abstract word count:** 244.

**Manuscript word count: 1956.**

Journal Pre-proof

**Abstract:**

The novel severe acute respiratory syndrome coronavirus-2 (SARSCoV-2) pandemic is a global health problem, which started to affect China by the end of year 2019. In Europe, Italy has faced this novel disease entity (named COVID-19) first and severely. COVID-19 represents a significant hurdle for public health services and a potential harm for patients with cancer. The Collegio Italiano dei Primari Oncologi Medici (CIPOMO) is an Italian association of head physicians in oncology departments, which promotes working and research activities in oncology on a national basis. In the midst of the epidemic in Italy, the CIPOMO promoted a national survey aiming to evaluate the impact of COVID-19 on clinical activity of oncologists and the implementation of containment measures of COVID-19 diffusion. Overall, 122 head physicians participated in this survey, with a homogeneous distribution on the national territory. Results show that the following measures for oncologic patients have been promptly implemented through the whole country: use of protective devices, triage of patients accessing the hospital, delay of non-urgent visits, and use of telemedicine. Results of this survey suggest that Italian oncology departments have promptly set a proactive approach to the actual emergency. Oncologists need to preserve the continuum of care of patients, as the benefit of ensuring a well-delivered anti-cancer treatment plan outweighs the risk of COVID-19 infection. International cooperation is an important starting point, as heavily affected nations can serve as an example to find out ways to safely preserve health activity during pandemic.

**Keywords:** cancer, infection, coronavirus, pandemic, health care, oncology.

**Acknowledgments:** We thank all the CIPOMO members for participating in the present survey, thereby making this work possible to be shared in the oncologists' community.

**Fundings:** no funding was obtained for the present investigation.

## Introduction

On March 11<sup>th</sup> 2020, the World Health Organization (WHO) declared the novel severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) outbreak a pandemic [1]. By the end of February 2020, Italy was experiencing the rapid spread of the virus, which started to affect the North of the Country with a daily increase in the number of cases and consequent deaths [2]. In Italy, data regarding the diffusion of the novel coronavirus disease (COVID-19), caused by SARS-CoV-2 confirmed its higher lethality compared to that observed in China and worldwide (9% vs 4.3%) [3].

Following the Chinese model, containment measures to reduce the risk of COVID-19 in Italy have been promptly activated and implemented. The first national decree, issued on March 8<sup>th</sup>, instituted a containment zone concerning the most affected areas of the Country (the so-called Red Zone, which at that time included 3 regions in the North of Italy: Lombardia, Emilia Romagna, and Veneto). In the following days, a series of decrees have extended increasingly strict measures to the whole national territory. The main provisions included: forbidding all gatherings of people; restricting movements of people within and outside the hometown, except for circumstances of necessity; encouraging employees to work from home. In this circumstance, health workers cannot take any leave, and are asked to suspend all non-urgent activities. All planned surgeries are postponed, to give over intensive care beds to the treatment of patients with COVID-19, and hospitals had to create new intensive care places by converting operating and anesthetic rooms. Table 1 outlines the key milestones of COVID-19 diffusion.

This pandemic represents a significant harm for patients with cancer, who are at high risk of infections due to several predisposing factors [4]. Moreover, most treatment procedures in oncology cannot be delayed without compromising the efficacy of treatment itself. In Italy, specific indications on oncologic patients' management were given on March 10<sup>th</sup>, mainly regarding caution measures to reduce the risk of infection (i.e. use of personal protective equipment, practice social distancing). While encouraging physicians to postpone follow up visits, indications were given to

guarantee oncologic services even within the Red Zone, in order to maintain the continuum of care. At the present time, data regarding diffusion and management of COVID-19 in oncologic patients are scarce [5,6]. Such emergency has led Italian oncologists to join forces, with the aim to find a way not to compromise the continuum of care of patients and to preserve safe everyday clinical practice.

The Italian College of Directors of the National Health System (NHS) Hospital Departments of Medical Oncology (CIPOMO - Collegio Italiano dei Primari Oncologi Medici Ospedalieri) gathers the heads of all the NHS hospital medical oncology departments, including general hospitals, city hospitals, and cancer centers. This association safeguards and promotes working and research activities in oncology on a national basis. In the midst of the epidemic in Italy, the CIPOMO promoted a national survey aiming to evaluate the impact of COVID-19 on clinical activity of oncologists and the implementation of containment measures of COVID-19 diffusion. Here we present the results of this survey, providing an overview of COVID-19 epidemic in Italy and assessing potential interventions to overcome this critical situation.

### **Survey characteristics and results**

The survey includes a total of 27 questions, which are divided in 3 sections: the first section assesses the routine use of preventive measures (e.g. vaccinations) in oncologic patients; the second contains questions regarding COVID-19 diffusion containment measures adopted before the enactment of national decrees in this regard; the third and last section assesses the diffusion of COVID-19 in oncology units and its impact on working activity, after national decrees on containment measures were adopted. A complete original version of the survey is provided in Supplementary Material 1.

The survey was launched online on March 12<sup>th</sup> 2020, and closed on March 15<sup>th</sup> 2020. Figure 1 displays the COVID-19 diffusion in Italy at the beginning and end of the present survey. At the time of the survey initiation, most of COVID-19 cases were gathered in the Red Zone. Overall, 122

head physicians of oncology departments participated in the survey, with an 84% survey completion rate. Results of the survey sections are displayed in Tables 2-4. In Italy there is a universalistic NHS, with an individual administration further subdivided among 20 regions which are geographically distributed in North, Centre, and South (including the islands). For this reason, results are presented as follows: all regions (n = 122); Red Zone (n = 39); North Italy, excluding the Red Zone (n = 26); Centre Italy (n = 18); South Italy and islands (n = 25). To simplify consultation and interpretation of results, Tables 2-4 contain answers to multiple choice questions of the survey, while answers to open questions are reported further in the main text.

Results of Section 1 show that the overall tendency throughout Italy is to perform preventive vaccination in oncologic patients (Table 2). Most vaccinated patients are either receiving active systemic treatment (chemotherapy, immune-therapy, and hormonal or targeted therapy), and/or present one or more predisposing risk factors (i.e. age  $\geq$  75 years, cardiovascular and/or respiratory disease, chronic infections, diabetes, obesity, immune-suppressive therapies). The most widely adopted vaccination is for seasonal flu, however more than 30% of oncologists suggests also performing pneumococcal vaccination in those patients. Seasonal flu cases are not usually reported by medical oncologists to the dedicated national registry, rather this is commonly a duty of family doctors.

Answers to the questions in Section 2 clearly show that, even if by the time of the survey COVID-19 represented an emergency mainly in the North of Italy, diagnostic measures for all patients accessing oncologic services were immediately activated in the whole country (Table 3). Triage of patients included vital signs monitoring before entering the hospital (body temperature, SpO<sub>2</sub>, respiratory rate), but also questioning patients on the presence of symptoms during the 15 days before the visit, and possible contacts with subjects affected by COVID-19 or coming from high-risk areas. In more than 65% of cases, triage procedure was followed by preventive isolation and diagnostic work up of symptomatic patients, consisting in chest X-ray and rhino-pharyngeal swab to rule out the presence of SARSCoV-2.



After the very first reports of COVID-19 in Italy, measures to reduce hospital accesses for oncologic patients were taken almost throughout the country. Such measures consisted mainly in delaying visits not considered to be urgent (i.e. patients in follow up after surgery and/or radiotherapy, and/or patients with breast cancer receiving adjuvant hormonal treatment after surgery), even more so if the patients presented risk factors (i.e. age > 85 years, presence of comorbidities). Alternative ways to get in touch with patients have been widely used: most patients underwent telephonic interviews with interpretation of laboratory and radiologic exams report, while in other cases family doctors were delegated to inspect the results of follow up exams. Access to oncologic structures was likewise limited and/or denied for visitors and caregivers, either for outpatient visits, day hospital and ward admissions.

With the enactment of the decree on March 9<sup>th</sup> and subsequent decree on March 11<sup>th</sup>, containment preventive measures became effective in the Red Zone before, and on the whole Italian territory thereafter. However, by that time most oncologic units had already activated measures to contain accesses, under regional or internal (hospital Medical Direction) orders. As so, more than 50% of oncologic structures did not have to modify the measures they implemented to reduce the risk of infections, in view of the actual legislation. Due to such measures, more than 20% of structures had reported a significant reduction in their routine activity, while 60% had only a negligible reduction and 9% had not substantially changed their activity.

Answers to the questions in Section 3 show that almost one third of oncologic structures had to employ their oncologists for guard duties in Internal Medicine ward and/or Emergency Department; in 23% of cases, guard duties in COVID ward were included (Table 4). This percentage was understandably higher in the North of the Country, reaching 51% and 38% of oncologists employed for Internal Medicine/Emergency Department and COVID wards guard duties, respectively, in the Red Zone. Twenty-four percent of Italian oncology departments had at least one patient diagnosed with COVID-19, with a higher rate in the Red Zone (46%) and no diagnosis at all in the South of Italy and in islands. Examining reports from the Red Zone, it

emerges that most patients accessed Emergency Room presenting with fever and/or respiratory symptoms (23%), while a significant proportion of patients was diagnosed after a triage procedure (18%) and/or a medical interview regarding possible contacts with subjects at risk (18%).

When questioned about the impact of containment measures in oncologic structures, most physicians found that measures could have a significant impact on reducing the risk of COVID-19 diffusion (53%), while 38% found they could have only a negligible effect. Answers to an open question asking whether other useful measures were applied (i.e. other than those indicated by decrees), raised the following issues: re-distribution of patients' treatment sessions homogeneously throughout the week; reduction of hospital accesses by relying on family doctors for home visits (e.g. medications, patients on treatment with oral metronomic chemotherapy); creation of a dedicated telephone line for emergencies; careful evaluation of risk-to-benefit ratio in heavily pre-treated patients, possibly postponing the start of further lines of palliative therapies. Regarding working activity, multidisciplinary boards have been converted in telematic meetings, and counseling of patients in other hospital wards have been managed by phone, when feasible. In some cases, physicians on duty in COVID wards have been consequently waived from oncologic activities, in order to reduce the risk of infection for both patients and colleagues.

## **Discussion**

Results of this survey deserve several considerations. Some effective points emerge from this survey, which partly have been also raised in the comment recently released by the European Society for Medical Oncology (ESMO) [7]. First, oncologists face the need to preserve the continuum of care for most of their patients, as the benefit of ensuring a well-delivered anti-cancer treatment plan outweighs the risk of COVID-19 infection. However, the risk from COVID-19 exposure varies significantly from patient to patient, making treatment tailoring important now more than ever. Second, physicians have to get used to a new working activity, which implies the use of tele-consultation services when feasible, and reducing the number of visits by means of

customizing treatment delivery (three or two-weekly as opposed to weekly, oral or subcutaneous alternatives as opposed to intravenous administration). Also, in view of a visit a telephonic “previous day” triage could help, in order to avoid the access of symptomatic patients to oncologic wards. Third, protection of patients and physicians is paramount in order to keep providing the best service in a safe way.

This last issue raises the question on how to optimize oncologic resources for COVID-19 emergency. Due to both clinical characteristics of oncologic patients, which are frail and at high risk for infections, and features of oncologic services, which often cannot be postponed, how can oncologist reasonably help in this emergency without compromising patients’ continuum of care? To date, no clear indications have been provided to health care providers in oncology, making it difficult to create a common line of action.

## **Conclusions**

In conclusion, results of our survey suggest that Italian oncology departments have promptly set a proactive approach to the actual emergency. Together with indications provided by national decrees and internal dispositions, several individual initiatives have contributed to reorganize working activity in the actual condition. The medical community worldwide is facing a difficult situation, and oncologists in particular require several extra precautions to protect the patients first and their activity thereafter. International cooperation is an important starting point, as heavily affected nations can serve as an example to find out ways to safely preserve health activity during pandemic.

**Conflicts of Interest statement**

The authors they have no conflicts of interest to declare for the present paper.

Journal Pre-proof

## References

- [1] The World Health Organization (WHO) website, available at <https://www.who.int/emergencies/diseases/novel-coronavirus-2019>, accessed March 22<sup>nd</sup> 2020.
- [2] Lazzerini M, and Putoto G. COVID-19 in Italy: momentous decisions and many uncertainties. *Lancet Glob Health*. 2020 Mar 18. doi: 10.1016/S2214-109X(20)30110-8. [Epub ahead of print].
- [3] The World Health Organization (WHO) Health Emergency Dashboard, available at <https://experience.arcgis.com/experience/685d0ace521648f8a5beeeee1b9125cd>, accessed March 19<sup>th</sup> 2020.
- [4] El Ramahi R, and Freifeld A. Epidemiology, Diagnosis, Treatment, and Prevention of Influenza Infection in Oncology Patients. *J Oncol Pract*. 2019; 15(4): 177-184.
- [5] Liang W, Guan W, Chen R, et al. Cancer patients in SARS-CoV-2 infection: a nationwide analysis in China. *Lancet Oncol*. 2020; 21(3): 335-337.
- [6] Yu J, Ouyang W, Chua MLK, et al. SARS-CoV-2 Transmission in Patients With Cancer at a Tertiary Care Hospital in Wuhan, China. *JAMA Oncol*. Published online March 25<sup>th</sup> 2020. doi:10.1001/jamaoncol.2020.0980
- [7] European Society for Medical Oncology (ESMO) Newsroom, available at <https://www.esmo.org/newsroom/covid-19-and-cancer/supporting-oncology-professionals>, accessed March 22<sup>nd</sup> 2020.

**Table 1.** Timeline of the key stages of COVID-19 diffusion in Italy.

December 31 <sup>st</sup> 2019	The Municipal Health Services in Wuhan (China) report to the WHO a cluster of patients with pneumonia of unknown etiologic agent in the city of Wuhan, in the Chinese province of Hubei.
January 9 <sup>th</sup> 2020	Chinese authorities make a preliminary determination of a novel (or new) coronavirus (SARSCoV-2), as the causal agent of the severe acute respiratory syndrome, named COVID-19. Chinese investigators conduct gene sequencing of the virus, using an isolate from one positive patient sample, making diagnostic tests promptly available worldwide.
January 22 <sup>nd</sup> 2020	The Italian Ministry of Health sets up a task force to coordinate interventions on the Italian territory, together with international responsible institutions. A surveillance system for suspected cases is established.
January 30 <sup>th</sup> 2020	Two Chinese tourists hospitalized for respiratory tract infection, in Rome, are the first confirmed cases of COVID-19 detected in Italy. Regional Health Authorities implement measures to track contacts of the two subjects. All contacts resulted negative for COVID-19. Italian government decides to interrupt all air connections with China. The WHO declares COVID-19 diffusion in China a public health emergency.
January 31 <sup>st</sup> 2020	The Italian Council of Ministers declares national public health emergency condition.
February 21 <sup>st</sup> 2020	The Italian National Institute of Health confirms the first case of local transmission of COVID-19 infection in a 38-year-old patient in Codogno.
February 28 <sup>th</sup> 2020	The WHO raises the threat definition for COVID-19 epidemic at a “high level” of threatening for the global health.
March 8 <sup>th</sup> -9 <sup>th</sup> 2020	A national decree institutes a containment zone concerning the most affected areas of the Country, located in the North of Italy (Lombardia, Emilia Romagna, and Veneto). The main provisions include: forbidding all gatherings of people; restricting movements of people within and outside the hometown, except for circumstances of necessity; encouraging employees to work from home. Health workers cannot take any leave, and are asked to suspend all non-urgent activities; all planned surgeries are postponed, to give over intensive care beds to the treatment of patients with COVID-19.
March 10 <sup>th</sup> 2020	The Italian Ministry of Health issues recommendations for oncologic and onco-hematologic patients, providing protective measures for off-therapy patients and those receiving systemic treatment. Oncologists are required to postpone follow up visits, in order to reduce patients’ access to hospitals. Patients with thoracic tumors and those who underwent pulmonary resection are to be considered a subgroup of high risk patients.
March 11 <sup>th</sup> 2020	The Italian Council of Ministers urgently sets increasingly strict containment measures, to the whole national territory. The WHO declares the novel SARSCoV-2 outbreak a pandemic.
March 13 <sup>th</sup> 2020	Three Italian scientific associations (AIOM, CIPOMO and COMU) release an official document for the management of oncologic and onco-hematologic activities during COVID-19 pandemic. The WHO declares Europe is becoming the new epicenter of COVID-19 pandemic.
Abbreviations: AIOM, Associazione Italiana di Oncologia Medica; CIPOMO, Collegio Italiano dei Primari Oncologi Medici Ospedalieri; COMU, Collegio degli Oncologi Medici Universitari; COVID-19, novel coronavirus disease; SARSCoV-2, severe acute respiratory syndrome coronavirus-2; WHO, World Health Organization.	

**Table 2.** Survey section 1: general (i.e. non COVID-19) preventive measures for reducing the risk of infections in oncologic patients.

		All, n (%) n = 122	Red Zone <sup>a</sup> , n (%) n = 39 (32)	North, n (%) n = 26 (21)	Centre, n (%) n = 18 (15)	South and islands, n (%) n = 25 (20)
Vaccination of oncologic pts in view of a systemic therapy	NO	8 (7)	2 (5)	2 (8)	2 (11)	0
	YES	114 (93)	37 (95)	24 (92)	16 (89)	25 (100)
Which kind of oncologic treatment will the vaccinated pts receive*	CT	28 (25)	8 (20)	7 (27)	4 (22)	6 (24)
	ICI	12 (11)	4 (10)	4 (15)	1 (5)	1 (4)
	HT, TT	16 (14)	6 (15)	4 (15)	2 (11)	3 (12)
	None, follow up	8 (7)	3 (8)	1 (4)	0	2 (8)
	All pts	64 (57)	22 (56)	16 (61)	5 (28)	16 (64)
	Pts with risk factors <sup>b</sup>	42 (37)	14 (36)	7 (27)	8 (44)	7 (28)
Which kind of vaccination is/are advised*	Seasonal flu	72 (64)	27 (69)	17 (65)	11 (61)	12 (48)
	Pneumococcal	8 (7)	3 (8)	2 (8)	0	0
	Both	42 (37)	11 (28)	8 (31)	4 (22)	13 (52)
	Other (Tetanus, VZV)	1 (<1)	1 (<1)	0	0	0
Report of seasonal flu cases (e.g. national registry)	NO	94 (80)	28 (72)	21 (81)	16 (89)	23 (92)
	YES	15 (13)	8 (20)	3 (11)	1 (5)	2 (8)
	UNK	9 (7)	3 (8)	2 (8)	1 (5)	0
* questions with more than one possible answer.						
<sup>a</sup> Red Zone at the time of the survey includes the following Italian countries: Lombardia, Emilia Romagna, Veneto.						
<sup>b</sup> age $\geq$ 75 years, cardiovascular and/or respiratory disease, chronic infections, diabetes, obesity (BMI $\geq$ 30), immune-suppressive therapies.						
Abbreviations: BMI, body mass index; CT, chemotherapy; HT, hormonal therapy; ICI, immune-checkpoint inhibitors; TT, targeted therapy; UNK, unknown; VZV, varicella zoster virus.						

**Table 3.** Survey section 2: COVID-19 preventive measures.

		All, n (%) n = 122	Red Zone <sup>a</sup> , n (%) n = 39 (32)	North, n (%) n = 26 (21)	Centre, n (%) n = 18 (15)	South and islands, n (%) n = 25 (20)
Triage process <sup>c</sup> for pts accessing oncologic services	NO	6 (5)	2 (5)	1 (4)	1 (6)	2 (4)
	YES	112 (95)	37 (95)	25 (96)	17 (94)	23 (96)
Which kind of pts are triaged*	All	79 (70)	24 (61)	24 (92)	10 (56)	14 (56)
	Pts with fever	20 (19)	7 (18)	1 (4)	5 (28)	6 (24)
	Pts with RS <sup>d</sup>	14 (12)	5 (13)	1 (4)	4 (22)	4 (16)
	Pts with fever <sup>e</sup> + RS	8 (7)	8 (20)	0	4 (22)	5 (20)
Pts are questioned about fever <sup>e</sup> , RS, and/or contact with high-risk subjects	NO	5 (4)	1 (2)	1 (4)	2 (11)	1 (4)
	YES	113 (96)	38 (97)	25 (96)	16 (89)	24 (96)
Diagnostic work-up for pts with suspicious symptoms	NO	39 (33)	9 (23)	6 (23)	9 (50)	13 (52)
	YES	79 (67)	30 (77)	20 (77)	9 (50)	12 (48)
Measures for reducing pts accesses from first cases of COVID-19	NO	11 (9)	5 (13)	2 (8)	1 (6)	1 (4)
	YES	107 (91)	34 (87)	24 (92)	17 (94)	24 (96)
Which kind of pts are these measures addressed to*	all follow up visits (non urgent)	98 (92)	30 (77)	24 (92)	15 (83)	22 (88)
	follow up visits of high-risk pts <sup>b</sup>	33 (31)	9 (23)	7 (27)	7 (39)	6 (25)
	pts receiving adjuvant therapy	4 (4)	0	1 (4)	1 (6)	2 (8)
Alternative modalities to get in touch with pts (e.g. telephonic interview)	NO	23 (20)	10 (26)	3 (11)	4 (22)	4 (16)
	YES	88 (80)	25 (64)	21 (81)	14 (78)	22 (88)
Reduced access for pts coming from Red Zone, and/or reporting fever <sup>e</sup> and/or RS	NO	21 (18)	9 (23)	3 (11)	6 (33)	2 (8)
	No, Red Zone out of catchment area	18 (15)	2 (5)	4 (15)	4 (22)	6 (24)
	YES	77 (66)	28 (72)	19 (73)	8 (45)	17 (68)
Reduced access for visitors and/or caregivers of pts in hospital ward, day hospital, outpatient visits	NO	0	1 (3)	0	0	0
	Yes, limited access for one visitor per patient	59 (51)	23 (59)	10 (38)	9 (50)	14 (56)
	Yes, denied access for visitors	50 (43)	13 (33)	15 (58)	9 (50)	11 (44)
Autonomous management of measures	YES	50 (43)	17 (44)	11 (42)	7 (39)	12 (48)
	Medical direction measures	53 (46)	18 (46)	13 (50)	6 (33)	12 (48)
	Regional measures	13 (11)	4 (10)	2 (8)	5 (28)	1 (4)
Measures for reducing pts accesses in view of	NO, internal measures	61 (52)	22 (56)	12 (46)	9 (50)	16 (64)



actual legislation <sup>f</sup> *	already activated					
	YES	59 (51)	19 (49)	16 (61)	9 (50)	9 (36)
Which kind of pts are these measures addressed to*	all follow up visits (non urgent)	82 (92)	26 (67)	21(81)	12 (67)	19 (76)
	follow up visits of high-risk pts <sup>b</sup>	26 (29)	9 (23)	4 (15)	3 (23)	8 (32)
	pts receiving adjuvant therapy	3 (3)	0	1 (4)	1 (8)	2 (8)
Did your Country provide guidelines for oncologic pts management during COVID-19 emergency?	NO	33 (28)	11 (28)	7 (27)	6 (33)	7 (28)
	YES	73 (63)	26 (67)	17 (65)	8 (45)	16 (64)
	Only for hospital pts	10 (9)	2 (5)	2 (8)	4 (22)	2 (8)
How is your activity changing in view of actual legislation <sup>f</sup> ?	Significant decreased	28 (24)	9 (23)	6 (23)	7 (39)	5 (20)
	Negligible decreased	70 (60)	23 (59)	18 (69)	9 (50)	15 (60)
	Unchanged	10 (9)	5 (13)	1 (4)	1 (6)	2 (8)
	Increased	0	0	0	0	0
	UNK	8 (7)	2 (5)	1 (4)	1 (6)	3 (12)

\*questions with more than one possible answer.

<sup>a</sup> Red Zone at the time of the survey includes the following Italian countries: Lombardia, Emilia Romagna, Veneto.

<sup>b</sup> age  $\geq$  75 years, cardiovascular and/or respiratory disease, chronic infections, diabetes, obesity (BMI  $\geq$  30), immune-suppressive therapies.

<sup>c</sup> Patients triage includes: vital signs, and respiratory signs/symptoms evaluation.

<sup>d</sup> Definition of fever: TC  $\geq$  37.5 °C.

<sup>e</sup> Definition of respiratory symptoms: dyspnea, cough, rhinitis.


<sup>f</sup> Administrative order issued on 11st March (see text for complete details).

Abbreviations: BMI, body mass index; RS, respiratory symptoms; UNK, unknown.

**Table 4.** Survey section 3: overview of COVID-19 diffusion.

		All, n (%) n = 122	Red Zone <sup>a</sup> , n (%) n = 39 (32)	North, n (%) n = 26 (21)	Centre, n (%) n = 18 (15)	South and islands, n (%) n = 25 (20)
Have your oncologists been requested for guard duties in Internal Medicine and/or Emergency wards?	NO	75 (61)	19 (49)	15 (58)	15 (83)	24 (96)
	YES	39 (32)	20 (51)	11 (42)	3 (17)	1 (4)
Are guard duties including COVID wards?	NO	29 (24)	9 (23)	5 (19)	6 (33)	7 (28)
	YES	28 (23)	15 (38)	9 (35)	1 (5)	1 (4)
Has any of your patients been diagnosed with COVID?	NO	86 (70)	21 (54)	20 (77)	14 (78)	25 (100)
	YES	29 (24)	18 (46)	6 (23)	4 (22)	0
How was COVID diagnosis performed?*	Contact with high risk subject(s) <sup>b</sup>	8 (6)	7 (18)	0	1 (5)	-
	Symptoms at triage	11 (9)	7 (18)	3 (8)	1 (5)	-
	Hospitalized patients and/or Emergency Room access	15 (12)	9 (23)	3 (8)	2 (11)	-
Do you reckon that actual measures will reduce the risk of COVID diffusion in oncologic wards?	NO	1 (<1)	1 (3)	0	0	0
	Yes, negligible	47 (38)	18 (46)	10 (38)	5 (28)	10 (40)
	Yes, significant	65 (53)	20 (52)	16 (62)	13 (72)	15 (60)
* questions with more than one possible answer.						
<sup>a</sup> Red Zone at the time of the survey includes the following Italian countries: Lombardia, Emilia Romagna, Veneto.						
<sup>b</sup> High risk subjects: people with known COVID infection and/or coming from Red Zone.						

Journal Pre-proof



Journal Pre-proof

**Legend to figures.**

**Figure 1.** Overview of COVID-19 diffusion in Italy at the beginning of the survey on March 12<sup>th</sup> 2020 (1a), and at the end of the survey on March 15<sup>th</sup> 2020 (1b).

Journal Pre-proof

**Highlights:**

- The novel coronavirus disease is a global health problem
- Patients with cancer are at risk of infections due to several predisposing factors
- Delaying oncologic procedures can compromise the efficacy of treatment itself
- Oncologists face the need to preserve their patients' continuum of care
- Proactive containment measures can protect both patients and medical activity

Journal Pre-proof

**Conflicts of Interest Statement**

All authors have no conflict of interest to disclose.

Journal Pre-proof