




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Available online at www.sciencedirect.com

ScienceDirect

journal homepage: www.e-jmii.com

Correspondence

Evacuation of quarantine-qualified nationals from Wuhan for COVID-19 outbreak – Taiwan experience

Dear Editor,

The 2019–20 coronavirus pandemic is an ongoing global pandemic of coronavirus disease 2019 (COVID-19) caused by the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). The virus was first reported in Wuhan, Hubei, China, in December 2019. After the lockdown of Wuhan and Hubei, the Taiwanese government planned to evacuate its citizens from the area by means of chartered flights. Out of the 47 laboratory-confirmed COVID-19 cases reported in Taiwan by March 10 2020, 21 were imported. A detailed evacuation workflow was planned in order to minimize the possibility of surface environmental contamination¹ - Pre-flight: (1) Equipment/facilities. (2) Regulation/communication. (3) Training/drills. Upon arrival at Wuhan airport: (1) Quarantine certification. (2) Self-report medical declaration. (3) Physical evaluation. Boarding: (1) Disinfection. (2) Protective clothing. (3) Boarding priority/seat arrangement. (Fig. 1) Inflight: (1) No inflight service. (2) Waste handling. (3) Monitor/seat-rearrangement. Upon arrival at Taiwan airport: (1) Disembark order. (2) Transport/self-health management.

On March 10 2020, Taiwan evacuated her second batch of citizens from Wuhan. A specialized medical team consisting of 4 physicians and 9 registered nurses was organized. Before flight, the medical team and their equipment were examined in detail. All the medical staff were equipped with personal protective gear (protective coveralls, face shield, N95 mask, gloves) and these remained donned throughout the mission.

At Wuhan airport, the evacuees were first screened by the mission nurse using an infrared thermometer. If their forehead temperature was 37.4 °C or more, it would be rechecked using an ear thermometer. If the second reading was still ≥ 37.4 °C, he/she was considered to be “febrile” and excluded from the evacuation.

Only afebrile evacuees would proceed to have their quarantine certification and self-report medical declaration verified by the medical team, following which they would be examined by the doctors. In addition to being febrile, other conditions that would lead to a denial of evacuation included (i) active respiratory symptoms with an oxygen saturation $\leq 95\%$ on pulse oximetry or abnormal respiratory examination findings, (ii) general physical condition that is not suitable for flight, and (iii) a pre-existing diagnosis of COVID-19 for which the evacuee may be receiving treatment, or still pending isolation or quarantine.

Those who were well on examination (afebrile with a pulse oximetry reading of $\geq 95\%$ and unremarkable pulmonary examination), and free of fever and respiratory symptoms for the preceding 14 days, were identified with a green sticker. Red stickers were used to identify evacuees who were well on examination, but had declared that they had fever or respiratory symptoms in the past 14 days. The third group of evacuees, identified by black stickers, consisted of those who were afebrile, but experienced any kind of respiratory symptoms at the point of examination.

After completing screening by the medical team, evacuees were instructed to disinfect both hands and wear personal protective equipment (protective gown, face shield, surgical mask and gloves) before being allowed to board the aircraft. The boarding sequence was based on their color labels. Evacuees with black stickers boarded first and were seated towards the back of the plane, from row 43 backwards. The next group to board was the red group, and finally the green. Two seats were left vacant between each passenger. During the flight, people were asked not to talk to each other, and not to consume food and drinks. Leaving their seats to go to the toilet was not allowed. Evacuees were advised to inform the medical staff if they experienced any discomfort. After landing in Taiwan, all evacuees were examined again. The throat swab samples were collected and underwent real-time polymerase-chain-reaction test by the local quarantine officers before being transported separately to designated

<https://doi.org/10.1016/j.jmii.2020.03.023>

1684-1182/Copyright © 2020, Taiwan Society of Microbiology. Published by Elsevier Taiwan LLC. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).



Figure 1. Boarding priority & seat arrangement (Airbus A330 – 300) – Evacuees with black stickers boarded first and were seated from row 43 backwards. The next was the red group from row 39 onwards, then the green group from row 7 backwards, and finally the medical staff from row 6 onwards.

dormitories for 14 days of self-quarantine. All of 361 evacuees were proved to be negative for SARS-CoV-2 infection later.

While the risk remains that asymptomatic evacuees may transmit the disease,² countries may still be obliged to evacuate these citizens from epidemic areas that have been locked down. We hope to share certain insights derived by the Taiwanese medical team during the execution of our evacuation. Our experience was limited to a short-haul flight – it takes around 3 h to travel from Wuhan to Taiwan. For flights of longer duration, there needs to be allowances made for food, beverage, and toileting needs.³ Indeed, given the restrictions imposed during the flight, certain high-risk populations, for example people who are prone to developing deep venous thrombosis or neuropsychiatric-related conditions,⁴ may require special precautions or medical attention. Another limitation was our workflow could not show preventive efficacy for environmental contamination or diseases transmission for SARS-CoV-2 infected evacuees. In addition, an isolation unit should be considered for in-flight emergencies or isolation.^{3,5}

References

1. Ong SWX, Tan YK, Chia PY, Lee TH, Ng OT, Wong MSY, et al. Air, surface environmental, and personal protective equipment contamination by severe acute respiratory syndrome Coronavirus (SARS-CoV-2) from a symptomatic patient. *Jama* 2020. <https://doi.org/10.1001/jama.2020.3227>.
2. Rothe C, Schunk M, Sothmann P, Bretzel G, Froeschl G, Wallrauch C, et al. Transmission of 2019-nCoV infection from an asymptomatic contact in Germany. *N Engl J Med* 2020;**382**: 970–1.
3. Gibbs SG, Herstein JJ, Le AB, Beam EL, Cieslak TJ, Lawler JV, et al. Review of literature for air medical evacuation high-level containment transport. *Air Med J* 2019;**38**:359–65.
4. Szilard I, Cserti A, Hoxha R, Gorbacheva O, O'Rourke T. International organization for migration: experience on the need for

medical evacuation of refugees during the Kosovo crisis in 1999. *Croat Med J* 2002;**43**:195–8.

5. Tsai SH, Tsang CM, Wu HR, Lu LH, Pai YC, Olsen M, et al. Transporting patient with suspected SARS. *Emerg Infect Dis* 2004;**10**:1325–6.

Chien-Yu Lee*

Department of Pediatrics, Tao-Yuan General Hospital,
Ministry of Health and Welfare, Tao-Yuan, Taiwan, ROC

Pi-Sheng Wang

Hospital and Social Welfare Organizations Administration
Commission, Ministry of Health and Welfare, Nantou,
Taiwan, ROC

Yuan-Der Huang

Department of Obstetrics and Gynecology, Chiayi Hospital,
Ministry of Health and Welfare, Chiayi, Taiwan, ROC

Yung-Ching Lin

Taiwan Centers for Disease Control, Ministry of Health and
Welfare, Taipei, Taiwan, ROC

Yung-Nien Hsu

Department of Family Medicine, Tao-Yuan General
Hospital, Ministry of Health and Welfare, Tao-Yuan,
Taiwan, ROC

Shih-Chung Chen

Ministry of Health and Welfare, Taipei, Taiwan, ROC

*Corresponding author. Department of Pediatrics, Tao-Yuan
General Hospital, 1492 Jhongshan Road, Taoyuan District,
Tao-Yuan City 33004, Taiwan, ROC.
Fax: +886 3 3699721 ext. 3557.

E-mail address: chienyu1028@yahoo.com.tw (C.-Y. Lee)

21 March 2020

Available online ■ ■ ■