



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Journal Pre-proofs

Original article

Biological and Epidemiological Trends in the Prevalence and Mortality due to Outbreaks of Novel Coronavirus COVID-19

Sultan Ayoub Meo, Thamir Al-Khlaiwi, Adnan Mahmood Usmani, Anusha Sultan Meo, David C. Klonoff, Thanh D. Hoang

PII: S1018-3647(20)30151-8
DOI: <https://doi.org/10.1016/j.jksus.2020.04.004>
Reference: JKUSUS 1085

To appear in: *Journal of King Saud University - Science*

Received Date: 2 March 2020

Revised Date: 1 April 2020

Accepted Date: 5 April 2020

Please cite this article as: S.A. Meo, T. Al-Khlaiwi, A.M. Usmani, A.S. Meo, D.C. Klonoff, T.D. Hoang, Biological and Epidemiological Trends in the Prevalence and Mortality due to Outbreaks of Novel Coronavirus COVID-19, *Journal of King Saud University - Science* (2020), doi: <https://doi.org/10.1016/j.jksus.2020.04.004>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier B.V. on behalf of King Saud University.



Biological and Epidemiological Trends in the Prevalence and Mortality due to Outbreaks of Novel Coronavirus COVID-19

Sultan Ayoub Meo¹, Thamir Al-Khlaiwi², Adnan Mahmood Usmani³, Anusha Sultan Meo⁴, David C Klonoff, Thanh D. Hoang⁵

^{1,2}Department of Physiology, ²Strategic Centre for Diabetes Research, College of Medicine, King Saud University, Riyadh, Saudi Arabia.

³Army Medical College, National University of Health Sciences, Rawalpindi, Pakistan

⁴Diabetes Research Institute, Mills-Peninsula Medical Center (Sutter Health), San Mateo, California, United States of America

⁵Walter Reed National Medical Center, Bethesda, MD, USA

Key Words: Biological, Epidemiological trends, Coronavirus, COVID-19, Prevalence, Outbreak.

Running title: Biological and Epidemiological trends of COVID-19

Correspondence:

Prof. Sultan Ayoub Meo, MBBS, Ph.D

Professor and Consultant in Clinical Physiology, Department of Physiology, College of Medicine, King Saud University, PO Box 2925 Riyadh-11461. Kingdom of Saudi Arabia, Email: sultanmeo@hotmail.com.com

Abstract

The novel coronavirus (Covid-19) infection outbreak has posed a major threat to international health system and economy. This study is aimed at investigating the biological and epidemiological trends in the prevalence and mortality due to outbreaks of novel coronavirus (COVID-19) infections. The data on the global outbreak of COVID-19, were obtained from World Health Organization (WHO), Worldometer, Centers for Disease Control and Prevention (CDC), and research institutes. The information was also recorded from research documents published in global scientific journals indexed in Pub Med and Institute of

Scientific Information (ISI) Web of Science on the trends in the prevalence and mortality due to COVID-19 infection outbreaks. The results show rising trends in the transmission, prevalence and mortality rate due to coronavirus COVID-19. During the period of December 29, 2019 through March 31, 2020, it has infected 750890 people worldwide, resulting in 36405 deaths with a mortality rate of 4.84%. The infections were more frequent among male gender with above 60 years of age. The mean growth rate index for total number of cases from January 23 to March 31, 2020 was 1.20 and growth rate index for mortality rate was 1.12. There was a positive association between the prevalence and mortality rate ($R^2=0.996$). The novel coronavirus COVID-19 is highly contagious and has affected a large number of people worldwide. It is still spreading with mutable prevalence and mortality outbreak trends. The global health officials have taken priority measures to prevent the further outbreaks of this emerging pathogen across the globe. However, the rising number of cases and mortality risk estimates are demonstrating that enhanced public health mediations, good hygienic conditions, social distancing and movement limitations may control the COVID-19 epidemics. **Key Words:** Biological trends, Coronavirus, Covid-2019, 2019-nCoV, Prevalence, Outbreak.

1. Introduction

Viral infections are the most frequent infectious diseases and are common triggers for constituting major health and socio-economic harms (Meo et al., 2020). In late December 2019, an earlier named novel coronavirus (2019-nCoV), currently named as coronavirus (COVID-19) emerged from Wuhan, China, resulted in a challenging outbreaks in many regions in China and expanding globally. Human infections with coronavirus COVID-19 have raised great public health apprehension globally. In the third millennium, three major threatening infectious diseases outbreaks happened. Following the outbreak of SARS-CoV in 2002 (Zhong et al., 2003) and the MERS-CoV in 2012 (Nassar et al., 2018; Nassar et al., 2018), the novel Coronavirus “COVID-19” has emerged globally and has threatened the entire world. The coronaviruses are zoonotic infections, can be transmitted from animal to animal, animal to human, and human to human (Li et al., 2020; World Health Organization 2020).

The Covid-2019 first appeared in the last week of December 2019. The infection cases were originated from bats, snakes, seafood among people living in or visiting Wuhan, China and human-to-human transmission has also been confirmed (World Health Organization, 2020; Wang et al., 2020; Callaway and Cyranoski, 2020). The biological and epidemiological trends in the prevalence and mortality rate are changing daily. Initially, China bears the large burden of the diseases, whereas the incidence is gradually increasing in other countries mainly the Europe and United States of America. Despite recent efforts to understand the novel Coronavirus COVID-19, the science community, researchers and general population would like more information about the current biological and epidemiological situations. This study aimed at investigating the

mutable biological and epidemiological trends in the prevalence and mortality outbreaks of coronavirus COVID-19 infections.

2. Materials and Methods

This study was conducted in the Department of Physiology, College of Medicine, King Saud University, Riyadh, Saudi Arabia. We obtained data on biological and epidemiological trends, global number of confirmed cases, and number of deaths due to Coronavirus COVID-19 infections all around the world. The data were obtained from the World Health Organization (WHO), Centers for Disease Control and Prevention (CDC), and reports from various countries and their allied ministries to the WHO. We also obtained data from search engines including (Worldometer-Coronavirus). Moreover, we also reviewed the literature published in the Institute of Scientific Information (ISI) Web of Knowledge, Thomson Reuters, Pub-Med, Medline, and appropriate findings were recorded. We assessed the growth factor, a factor by which quantity multiplies itself over time; daily cases divided by cases on the previous day. A growth factor more than 1.0, indicates an increasing pattern, whereas values between 0-1.0 show a decline pattern (Worldometer-Coronavirus).

Ethical Statement In this study the information on the biological and epidemiological trends on global prevalence and mortality due to novel Coronavirus Covid-2019 infections was obtained from the World Health Organization, Worldometer-Coronavirus, the Centers for Disease Control and Prevention and reports from various countries and their allied ministries to the WHO; hence ethical approval was not required.

Statistical Analysis

The data were recorded and analyzed, and the results were expressed in numbers and percentages. The regression analysis was performed; and a p -value <0.05 was considered significant.

3. Results

These results are established from the available information at the time of publication originating from the mentioned sources. The data were interpreted with carefulness however, the outbreak is evolving rapidly and there is swift change in the numbers. The global number of cases and number of deaths due to novel coronavirus COVID-19 infection are presented in Table I, and Figures 1 and 2. Moreover, epidemiological trends were presented and correlations were established (Figure 2-4). The novel coronavirus COVID-2019 has infected people in 198 countries and territories.

The total number of cases worldwide are 750890 people and the disease has caused 36405 (4.84%) deaths during the period December 29, 2019 to March 31, 2020 (Table I). The majority of these coronavirus cases, were reported from European Region 423946 (56.45%), American Region 163014 (21.70%), Western Pacific Region 104868 (13.96%), Easter Mediterranean Region 50349 (6.70%), South-East Asia Region 4215

(0.56%) and African Region 3786 (0.50%) (Table 1). In these contents COVID-19 highly effected the various countries including United States of America, Italy, France, Spain, Germany, Switzerland, and Iran (Figure 1). However, the highest mortality rate was found in Italy, Spain, United Kingdom, France and Iran (Figure 5). The outbreak of novel coronavirus Covid-2019 started on December 29, 2019. The results shows rising trends in the transmission, prevalence and mortality rate due to coronavirus Covid-2019.

In this study, we also analyzed the daily cases and death growth factors of the coronavirus Covid-2019. It was identified that during the January 22 to March 31, 2020 the growth factor of the number of cases worldwide was 0.28 to 6.96 (Mean 1.20) and the growth factor of death rates was 0.51 to 2.17 (Mean 1.12). There was a positive association between the prevalence and mortality rate ($R^2=0.996$). It showed a mutable change in both the daily case growth factor and mortality rate (Figure 3).

4. Discussion

Coronavirus infection is an emerging global health concern and has infected a significant portion of the world's population. In this study, we investigated the biological and epidemiological trends in the prevalence and mortality due to outbreaks of novel coronavirus COVID-19 infection. The COVID-19 infection is expanding over 198 countries and territories, infected 750890 people, has caused 36405 (4.84%) deaths during the period December 29, 2019 to March 31, 2020 (Table I).

The coronavirus Covid-2019 infection is rapidly transmitted, the disease is still in its spreading phase with predictions of infection expand to over one people globally. We have derived the estimates of the ongoing COVID-19 epidemic and found that still the mean growth factor for number of cases and mortality rate is more than 1.0. It is highly contagious because of its biological characteristics (Meo et al., 2020). It is a zoonotic disease with person-to-person transmission through droplet or contact transmission (Wu et al., 2019). Infected people can transmit the disease before they present clinical symptoms (Rothe et al., 2020). This is the main reason that the disease could swiftly spread from Wuhan, China to various corners of the globe. More recently, the epicenter of the diseases appears to have shifted from Wuhan, China to the European world and mainly Italy, and United States of America (World Health Organization, 2020). Many countries including China, Italy, Saudi Arabia, New Zealand, Poland, Ireland, Denmark, Spain, France have cancelled international flights and implemented the world's largest and most restrictive quarantines policies to minimize the further transmission and disease burden. For the prevention of the spread of the disease, these countries are not permitting people to enter or leave the country without strict screening of the passenger for Covid-2019 infection at airports.

(Quilty et al., (2020) evaluated the usefulness of thermal passenger screening for COVID-19 infection at airports exit and entry points. The authors found that effectiveness of entry screening was dependent on the effectiveness of the exit screening at the destination. The authors identified that 46% of the infected travelers were not detected, because of the incubation period, and asymptomatic or subclinical cases (Chan et al., 2020). (Quilty et al., 2020) also reported that airport screening is unlikely to detect a sufficient proportion of COVID-19 infected travelers. The screening is intended to be a barrier for preventing infected people from entering into the country; however, evidence of its effectiveness remains limited. These are reasons that health officials are still unable to control the spread of this contagious disease. The transmission of the disease is closely linked to the asymptomatic and infected people and those visiting the countries with COVID-19 pandemic.

The coronavirus COVID-2019 spreads over 198 geographical regions of the globe. The gender-based analysis demonstrated that the identified cases consisted mostly of men with median age range of 50-65 years. (Li et al., 2020) reported that the novel coronavirus Covid-2019 infection was found in China, and later spread almost entire nations of the world. The virus infection mainly occurred among the elderly with the median age of 59 years. December 29, 2019 to March 31, 2020. The number of cases was positively linked to the death rate of COVID-19 infection (Figure 4). The disease is still spreading with mutable prevalence and mortality outbreak trends.

5. Conclusions

The epidemiological trends show that novel coronavirus COVID-19 is highly contagious and has affected 750890 people, with a mortality rate of 4.84%. The number of cases were proportional to the number of deaths. There was a great fluctuation in the biological and epidemiological trends both in the growth factor of number of cases and the mortality rates. Global health officials have taken high priority measures to prevent the further outbreaks of this emerging pathogen across the globe, but still the coronavirus COVID-19 is swiftly spreading with mutable biological trends. The rising number of cases and mortality risk estimates are demonstrating the dire need for enhanced public health mediations, good hygienic conditions, social distancing and movement limitations to control the COVID-19 epidemics.

Acknowledgements

Researchers Supporting Project Number (RSP-2019/47), King Saud University, Riyadh, Saudi Arabia.

Conflicts of interest

The authors declare that they have no conflict of interests.


Table I. Worldwide number of laboratories confirmed cases and deaths due to Coronavirus Covid-2019.

Number	Total	(%)	Number of Deaths	(%)
--------	-------	-----	------------------	-----

Total number of cases	750890	100	36405	4.84
Western Pacific Region	104868	13.96	3671	3.50
European Region	423946	56.45	26694	6.29
South-East Asia Region	4215	0.56	166	3.93
Easter Mediterranean Region	50349	6.70	2945	5.84
American Region	163014	21.70	2836	1.73
African Region	3786	0.50	77	2.0
Preexisting conditions	CAD 13.2%; Diabetes 9.2%; Respiratory diseases 8%			
Age group	60-69=3.6%; 70-79=8%; >80 21.9%			

Note: World Health Organization, Data updated on March 31, 2020


Figure 1. Worldwide prevalence of COVID-19 infection



Note: Novel Coronavirus "COVID-19" data is presented from Dec 29, 2019 to March 30, 2020.


Countries with less than 2000 cases are not presented in this graph Ref: (Worldometer Coronavirus, 2020)

Figure 2. Trend of global prevalence of number of cases and deaths due to COVID-19 infection



Note: Novel coronavirus “COVID-19” data is presented from Jan 22, 2020 to March 31, 2020. Ref. (Worldometer-Coronavirus, 2020)

Figure 3. Growth factor of number of cases and deaths due to COVID-19 infection.



Note: “COVID-19” data is presented from Jan 22, 2019 to March 31, 2020. Ref: Worldometer- Coronavirus, 2020.

Figure 4. Correlation between number of cases and number of deaths due to COVID-19 infection.





Figure 5. Number of cases and deaths presented with 100% stacked graph



Note: COVID-19 data is presented from Jan 22, 2020 to March 31, 2020. (Worldometer-Coronavirus, 2020).

References

Callaway, E., Cyranoski, D., 2020. China coronavirus: six questions scientists are asking. *Nature*, 577: 605-607.

Centers for Disease Control and Prevention. Available at: <https://www.cdc.gov/> cited date March 16, 2020.

Chan, J.F.W., Yuan, S., Kok, K.H., To, K.K.W., Chu, H., Yang, J., 2020. A familial cluster of pneumonia associated with the 2019 novel coronavirus indicating person-to-person transmission: a study of a family cluster. *Lancet*. S0140-6736 (20)30154-9.

COVID-19 Coronavirus. Available at: <https://www.worldometers.info/coronavirus/> Cited date Feb 29, 2020.

ISI Web of Knowledge. Thomson Reuters, Available at: <http://webofknowledge.com/JCR/JCR?PointOfEntry=Home&SID=4FeKpokbnHkLlmE10Ge>, Cited date March 30, 2020.

Li, Q., Guan, X., Wu, P., Wang, X., Zhou, L., Tong, Y., 2020. Early transmission dynamics in Wuhan, China, of novel coronavirus-infected pneumonia. *N Engl J Med*. doi: 10.1056/NEJMoa2001316.

Meo, S.A., Alhowaikan, A., Al-khlaiwi, T., Meo, I.M., Halepoto, D.M., Iqbal, M., Usmani, A.M., Hajjar, W., Ahmed, N. 2020. Novel coronavirus 2019-nCoV: prevalence, biological and clinical characteristics comparison with SARS-CoV and MERS-CoV. *Eur Rev Med Pharmacol Sci*. 24 (4): 2012-2019.

Nassar, M.S., Bakhrebah, M.A., Meo, S.A., Alsuabeyl, M.S., Zaher, W.A. 2018. Middle East Respiratory Syndrome Coronavirus (MERS-CoV) infection: epidemiology, pathogenesis and clinical characteristics. *Eur Rev Med Pharmacol Sci*, 22: 4956-4961.

Nassar, M.S., Bakhrebah, M.A., Meo, S.A., Alsuabeyl, M.S., Zaher, W.A., 2018. Global seasonal occurrence of middle east respiratory syndrome coronavirus (MERS-CoV) infection. *Eur Rev Med Pharmacol Sci*, 22: 3913-3918.

Pub Med. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/?term=Coronavirus>. cited date March 30, 2020.

Quilty, B.J., Clifford, S., 2020. Cmmid nCoV Working Group, Flasche S, Eggo RM. Effectiveness of airport screening at detecting travellers infected with novel coronavirus (2019-nCoV). *Euro Surveill*. 25 (5). doi: 10.2807/1560-7917.ES.2020.25.5.2000080.

Rothe, C., Schunk, M., Sothmann, P., Bretzel, G., Froeschl, G., Wallrauch, C. 2020. Transmission of 2019-nCoV Infection from an Asymptomatic Contact in Germany. *N Engl J Med*. doi: 10.1056/NEJMc2001468.

Wang, W., Tang, J., Wei, F. 2020. Updated understanding of the outbreak of 2019 novel coronavirus (2019-nCoV) in Wuhan, China. *J Med Virol*. doi: 10.1002/jmv.25689.

World Health Organization. Available at: <https://www.who.int/emergencies/mers-cov/en/> cited date March 31, 2020.

World Health Organization: coronavirus. Available at: <https://www.who.int/health-topics/coronavirus>, cited date March 31, 2020.

Wu, Y.C., Chen, C.S., Chan, Y.J., 2020. Overview of the 2019 Novel Coronavirus (2019-nCoV): The Pathogen of Severe Specific Contagious Pneumonia (SSCP). *J Chin Med Assoc*. doi: 10.1097/JCMA.0000000000000270.

Zhong, N.S., Zheng, B.J., Li, Y.M., Poon, Xie, Z.H., Chan, K.H., Li, P.H., Tan, S.Y., Chang, Q., Xie, J.P., Liu, X.Q., Xu, J., Li, D.X., Yuen, K.Y., Peiris, Guan. Y. 2003. Epidemiology and cause of severe acute respiratory syndrome (SARS) in Guangdong, People's Republic of China in February 2003. *Lancet* 362: 1353-1358.

The Honorable Editor

Journal of King Saud University-Science

Sub: Biological and Epidemiological Trends in the Prevalence and Mortality due to Outbreaks of Novel Coronavirus COVID-19

I state that the manuscript has been submitted solely to Journal of King Saud University-Science and that it has not been previously published, either in whole or in part, nor have the findings been posted online. I also take responsibility to all aspects of the research and writing process, and take final responsibility for the paper. We declare no conflicts of interests.

Thank you and best regards,

Prof. Sultan Ayoub Meo, MBBS, Ph.D,

Professor and Consultant in Clinical Physiology, Department of Physiology, College of Medicine, King Saud University, PO Box 2925 Riyadh-11461. Kingdom of Saudi Arabia, Email: sultanmeo@hotmail.com / smeo@ksu.edu.sa

Journal Pre-proofs