



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

available at [www.sciencedirect.com](http://www.sciencedirect.com)  
journal homepage: [www.europeanurology.com](http://www.europeanurology.com)



European Association of Urology



## Research Letter

# Severe Acute Respiratory Syndrome Coronavirus 2 Infection in Renal Failure Patients: A Potential Covert Source of Infection

Yu Xiao<sup>a</sup>, Kaiyu Qian<sup>a,b</sup>, Yongwen Luo<sup>a</sup>, Song Chen<sup>a</sup>, Mengxin Lu<sup>a</sup>, Gang Wang<sup>b</sup>, Lingao Ju<sup>b</sup>, Xinghuan Wang<sup>a,b,\*</sup>

COVID-19, a highly infective disease caused by a newly identified coronavirus, severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2; previously 2019-nCoV), is spreading around the world [1]. Increasing evidence has confirmed the human-to-human transmission. A special group of COVID-19 patients is comorbid with chronic kidney disease (CKD) [2]. In patients with CKD, innate and adaptive immune function impairment would result in increased susceptibility to bacterial and viral infections. Therefore, SARS-CoV-2 infection of these dialysis patients has attracted our attention.

### High risk

As reported in our previous study, 41.30% of COVID-19 patients were likely infected in hospital. Most renal failure patients were immunocompromised due to uremia and needed routine dialysis in hospital two to three times per week. Therefore, renal failure patients should be more susceptible to SARS-CoV-2 than the normal population. We noticed that, as chest computed tomography scans show, approximately 20–30% of dialysis patients were diagnosed with suspected infection in multiple dialysis centers in the early period of COVID-19 outbreak. Intriguingly, most of these suspected cases have no obvious clinical symptoms, including fever, fatigue, and dry cough. However, these asymptomatic patients may be a potential source of infection, transmitting the virus to healthcare providers, their family members, and other patients. Therefore, SARS-CoV-2 infection in dialysis patients should be taken more seriously due to a high risk of asymptomatic transmission.

### Diagnosis

Healthcare-associated pneumonia is a common infectious problem encountered in hemodialysis patients [3,4]. Moreover, many dialysis patients suffer concurrently from

pulmonary edema, which may mimic pneumonia, in terms of both presenting with an abnormal chest radiograph and producing a similar clinical symptom. Nucleic acid test could provide a valuable support, but the sensitivity of the current nucleic acid tests needs to be improved. Given all of the above, the precise diagnosis of COVID-19 infection is more difficult in dialysis patients than in the normal population.

### Treatment

Another concern is the pharmacokinetics in these renal failure patients. Many antiviral drugs are eliminated by hepatic metabolism and renal excretion. There are few studies on the pharmacokinetics of antiviral drugs in patients with renal dysfunction. In addition, hemodialysis could take away some drugs and reduce their concentration in the blood. Taken together, an adjusted or even supplemental dose of those antiviral drugs should be administered.

In addition, a previous study has indicated that renal dysfunction in patients with pneumonia is associated with a risk of severe infection [5]. A higher mortality rate should be observed in renal failure patients infected with the coronavirus. Indeed, SARS-CoV-2 infection can induce acute renal injury, and this might superimpose the patient with an underlying renal problem [2].

In conclusion, SARS-CoV-2 infection should be monitored intensively for dialysis patients due to immunosuppression, difficulty in diagnosis, and additional concern regarding the use of antiviral drugs. Our report should prompt experts and medical workers to pay special attention to the SARS-CoV-2 infection of dialysis patients.

**Funding support:** This work was supported by the grant from the Medical Science Advancement Program of Wuhan University (TFLC2018002).

<https://doi.org/10.1016/j.eururo.2020.03.025>

0302-2838/© 2020 European Association of Urology. Published by Elsevier B.V. All rights reserved.

**Conflicts of interest:** The authors have nothing to disclose.

## References

- [1] Li Q, Guan X, Wu P, et al. Early transmission dynamics in Wuhan, China, of novel coronavirus-infected pneumonia. *N Engl J Med*. In press. <https://doi.org/10.1056/NEJMoa2001316>.
- [2] Wang D, Hu B, Hu C, et al. Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus-infected pneumonia in Wuhan, China. *JAMA*. In press. <https://doi.org/10.1001/jama.2020.1585>.
- [3] Carratala J, Garcia-Vidal C. What is healthcare-associated pneumonia and how is it managed? *Curr Opin Infect Dis* 2008;21:168–73.
- [4] Betjes MG. Immune cell dysfunction and inflammation in end-stage renal disease. *Nat Rev Nephrol* 2013;9:255–65.
- [5] Al-Tawfiq JA, Hinedi K, Ghandour J, et al. Middle East respiratory syndrome coronavirus: a case-control study of hospitalized patients. *Clin Infect Dis* 2014;59:160–5.

<sup>a</sup>Department of Urology, Zhongnan Hospital of Wuhan University, Wuhan, China

<sup>b</sup>Cancer Precision Diagnosis and Treatment and Translational Medicine, Hubei Engineering Research Center, Zhongnan Hospital of Wuhan University, Wuhan, China

\*Corresponding author. Department of Urology, Zhongnan Hospital of Wuhan University, Wuhan 430071, China. Tel. +86 2767813104. E-mail address: [wangxinghuan@whu.edu.cn](mailto:wangxinghuan@whu.edu.cn) (X. Wang).