

1 **Title Page**

2

3 **Forecasting the Cumulative Number of COVID-19 Deaths in China: a**
4 **Boltzmann Function-based Modeling Study**

5 Yuanyuan Gao ^{1, #}, Zhongyan Li ^{1, #}, Qi Ying ², Cheng Long ^{3, *} and Xinmiao Fu ^{1, *}

6

7 ¹ Provincial University Key Laboratory of Cellular Stress Response and Metabolic Regulation, College of Life
8 Sciences, Fujian Normal University, Fuzhou City, Fujian Province 350117, China

9 ² Department of Civil and Environmental Engineering, Texas A&M University, College Station, TX 77843,
10 USA.

11 ³ Department of Orthopaedic, Sichuan University West China Hospital, Chengdu City, Sichuan Province
12 610041, China

13

14 **Running title:** Forecasting COVID-19 Deaths in China

15

16 # These authors contributed to this work equally.

17 * To whom correspondence should be addressed to Professor Xinmiao Fu (xmfu@fjnu.edu.cn)
18 and Dr. Cheng Long (longchenghx@126.com)

19

20 **Keywords:** COVID-19, SARS-CoV-2, 2019-nCoV, Boltzmann function, coronavirus, SARS,
21 epidemic

22 **Word count:** 1020

23

24 **Abstract**

25 **The COVID-19 outbreak is on-going in China. Here we estimated the potential total**
26 **numbers of COVID-19 deaths in China, outside Hubei (in China), Hubei Province, Wuhan**
27 **City and outside Wuhan (in Hubei) by Boltzmann function-based analyses, which are 3342**
28 **(95% CI, 3214, 3527), 111 (109, 114), 3245 (3100, 3423), 2613 (2498, 2767) and 627 (603, 654),**
29 **respectively. The results may help to evaluate the severity of COVID-19 outbreaks and**
30 **facilitate timely mental service for the families of passed patients.**

31
32 An outbreak of 2019 novel coronavirus diseases (COVID-19) caused by SARS-CoV-2 is on-
33 going in China and has spread worldwide ^[1, 2]. As of Feb 13, 2020, there have been over 80000
34 confirmed COVID-19 patients and over 3000 deaths in China, most of which are in the epicenter
35 of the outbreak, Wuhan city and related regions in Hubei province. Although the number of new
36 confirmed cases has substantially decreased since Feb 13, 2020 and the outbreak appears to
37 approach the late phase in China, people have raised grave concerns about the severity of the
38 outbreak, especially questioning how many patients will die eventually. Here we estimated the
39 potential total number of COVID-19 deaths by applying Boltzmann function-based regression
40 analysis, an approach we recently developed for estimating the potential total numbers of
41 confirmed cases for both the ongoing SARS-CoV-2 outbreak and the gone 2003 SARS epidemic
42 ^[3].

43
44 We collected data for analysis on the officially released cumulative numbers of deaths in mainland
45 China, other provinces than Hubei, Hubei Province, Wuhan City, and other cities in Hubei (from
46 Jan 21 to Feb 29, 2020). We first verified that the cumulative numbers of confirmed cases with
47 respect to each region were all well fitted to the Boltzmann function (R^2 all being close to 0.999);
48 **Fig. 1A**), consistent with our earlier report using the data from Jan 21 to Feb 14, 2020 ^[3].
49 Assuming that the number of deaths is proportional to the number of confirmed cases for the

50 outbreak under specific circumstances, we speculated that the cumulative number of COVID-19
51 deaths would also obey the Boltzmann function. In support of this speculation, the cumulative
52 numbers of COVID-19 deaths in the above regions were all well-fitted to the Boltzmann function
53 (R^2 all being close to 0.999; **Figs. 1B, 1C** and **Table 1**), with the potential total numbers of deaths
54 being estimated as 3200 ± 40 , 108 ± 1 , 3100 ± 40 , 2500 ± 40 and 604 ± 6 respectively (**Table 1**). This
55 result, in conjunction with our earlier observation that the cumulative numbers of confirmed cases
56 of 2003 SARS in mainland China and worldwide were well fitted with the Boltzmann function,
57 prompted us to analyze the cumulative numbers of 2003 SARS deaths in the same way.
58 Consistently, we observed that the cumulative numbers of 2003 SARS deaths in mainland China,
59 Hong Kong and worldwide were all well fitted to the Boltzmann function (**Fig. 1D**), strongly
60 suggesting that the Boltzmann function is suitable to simulate the course of deaths associated with
61 coronavirus-caused diseases.

62
63 One issue regarding our analyses is that some COVID-19 deaths might be miss-reported such that
64 the reported death numbers represent a lower limit. For instance, 134 new deaths were suddenly
65 counted from more than 13000 clinically diagnosed patients in Hubei Province on Feb 12, 2020
66 (as clearly indicated by a sudden jump of deaths in **Fig. 1B**). Another uncertainty might result
67 from those unidentified COVID-19 deaths at the early phase of the outbreak. We applied the
68 Monte Carlo method (for detail, refer to the Methods section in SI file) to estimate such
69 uncertainty assuming that the relative uncertainty of the reported numbers of deaths follows a
70 single-sided normal distribution with a mean of 1.0 and a standard deviation of 2.5%. The
71 potential total numbers of COVID-19 deaths in the above regions were estimated to be 3260 (95%
72 CI 3187, 3394), 110 (109, 112), 3174 (3095, 3270), 2550 (2494, 2621) and 617 (607, 632),
73 respectively (**Figs. 1E** and **S1**), which are slightly higher than those estimated without uncertainty
74 (refer to **Table 1**).

75

76 To verify our Boltzmann function-based estimations, we calculated the potential total numbers of
77 deaths in the above regions by applying Richards function-based regression analyses, which had
78 been explored to simulate the cumulative numbers of confirmed cases of 2003 SARS in different
79 regions ^[4,5]. The potential total numbers of COVID-19 deaths in mainland China, other provinces,
80 Hubei Province, Wuhan City and other cities were estimated to be 3342 (3214, 3527), 111 (109,
81 114), 3245 (3100, 3423), 2613 (2498, 2767) and 627 (603, 654), respectively (**Fig. S2**), which are
82 close to what are estimated by the Boltzmann function-based analyses (**Table 1**). In addition, we
83 found that the established Boltzmann function was able to predict the death course in a short
84 period such that the released cumulative numbers of deaths from Mar 1 to 5 Mar 5 are close to
85 the estimated numbers, as exemplified in Wuhan (**Fig. 1F** and **Table S1**). If the data from Jan 21
86 to different closing dates were arbitrarily analyzed, we found that the course of COVID-19 deaths
87 could be largely simulated based on the data as of Feb 14 (**Fig. 1F**), implicating that the
88 Boltzmann function-based analysis could predict the trend ahead of approximate three weeks.

89
90 Collectively, we observed that all sets of data from both the COVID-19 deaths and the 2003
91 SARS deaths were well fitted to the Boltzmann function. We propose that the Boltzmann function
92 is suitable for analyzing not only the cumulative number of confirmed COVID-19 cases, as
93 reported by us recently ^[3] (also refer to **Fig. 1A**), but also those of deaths as reported here. We
94 noticed that modeling studies on the COVID-19 outbreak have been performed ^[6] and COVID-
95 19 deaths have been estimated by other groups using different models. For instance, Li et al
96 recently showed ^[7] by data driven analysis that a total of deaths in Hubei would be 2250, a number
97 much lower than the observed (2761 as of Feb 29). Using the Susceptible-Infected-Recovered-
98 Dead model Anastassopoulou et al forecasted that the total death might exceed 7000 by Feb 29
99 ^[8], a number apparently much higher.

100
101 Since case fatality ratio in the epicenter of the outbreak is still much higher than that in other

102 provinces of mainland China ^[1, 2], there is a great potential for government to optimize
103 preparedness and medical resource supplies therein, by which hundreds of lives of COVID-19
104 patients, particularly those severe and critically ill patients ^[2, 9], might be saved. In addition, our
105 estimates on the course of COVID-19 deaths (refer to **Table S2**) may benefit the mental health
106 service that needs to be timely provided to the families of passed patients ^[10].

107

108 **Acknowledgments**

109 This work is support by the National Natural Science Foundation of China (No. 31972918 and
110 31770830 to XF).

111 **Declaration of interest statement:** All authors declare no conflicts of interests.

112 **References**

- 113 1. Report of the WHO-China Joint Mission on Coronavirus Disease 2019 (COVID-19). 16-24 February 2020.
114 p. [https://www.who.int/docs/default-source/coronaviruse/who-china-joint-mission-on-covid-19-final-](https://www.who.int/docs/default-source/coronaviruse/who-china-joint-mission-on-covid-19-final-report.pdf)
115 [report.pdf](https://www.who.int/docs/default-source/coronaviruse/who-china-joint-mission-on-covid-19-final-report.pdf).
- 116 2. Epidemiology Working Group for NCIP epidemic Response, Chinese Center for Disease Control and
117 Prevention. The epidemiological characteristics of an outbreak of 2019 novel coronavirus diseases (COVID-
118 19) in China. *Chin J Epidemiol*, 2020. 41(2) 145-151.
- 119 3. Fu, X., et al., Simulating and Forecasting the Cumulative Confirmed Cases of SARS-CoV-2 in China by
120 Boltzmann Function-based Regression Analyses. *Journal of Infection*, 2020;
121 <https://doi.org/10.1016/j.jinf.2020.02.019>.
- 122 4. Hsieh, Y.H., Lee, J.Y., and Chang, H.L., SARS epidemiology modeling. *Emerging Infect Dis*, 2004. 10(6):
123 1165–1167.
- 124 5. Wang, X.S., Wu, J., and Yang, Y., Richards model revisited: Validation by and application to infection
125 dynamics. *Journal of Theoretical Biology*, 2012. 313 12–19.
- 126 6. Wu, J.T., Leung, K., and Leung, G.M., Nowcasting and forecasting the potential domestic and international
127 spread of the 2019-nCoV outbreak originating in Wuhan, China: a modelling study. *Lancet*, 2020; DOI:
128 10.1016/S0140-6736(20)30260-9.
- 129 7. Li, Q., Feng, W., and Quan, Y.H., Trend and forecasting of the COVID-19 outbreak in China. *Journal of*
130 *Infection*, 2020; <https://doi.org/10.1016/j.jinf.2020.02.014>.
- 131 8. Anastassopoulou, C.e.a., DATA-BASED ANALYSIS, MODELLING AND FORECASTING OF THE
132 NOVEL CORONAVIRUS (2019-NCOV) OUTBREAK. <https://doi.org/10.1101/2020.02.11.20022186>,
133 2020.
- 134 9. Yang, X., et al., Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in
135 Wuhan, China: a single-centered, retrospective, observational study. *Lancet Respir Med*, 2020;
136 [https://doi.org/10.1016/S2213-2600\(20\)30079-5](https://doi.org/10.1016/S2213-2600(20)30079-5).
- 137 10. Xiang, Y.T., et al., Timely mental health care for the 2019 novel coronavirus outbreak is urgently needed.
138 *Lancet Psychiatry*, 2020. 7(3) 228-229.

139

140

141

142

143
144
145
146
147
148


149 **Table 1 Summary of the estimated total numbers of COVID-19 deaths in China**

| Regions | Boltzmann ^a | | Boltzmann ^a | Richards ^a |
|-----------------------|------------------------|----------------|------------------------|-----------------------|
| | Mean±SD | R ² | (mean, 95% CI) | (mean, 95% CI) |
| mainland China | 3200±40 | 0.999 | 3260 (3187, 3394) | 3342 (3214, 3527) |
| other provinces | 108±1 | 0.996 | 110 (109, 112) | 111 (109, 114) |
| Hubei Province | 3100±40 | 0.999 | 3174 (3095, 3270) | 3245 (3100, 3423) |
| Wuhan City | 2490±40 | 0.998 | 2550 (2494, 2621) | 2613 (2498, 2767) |
| other cities in Hubei | 604±6 | 0.999 | 617 (607, 632) | 627 (603, 654) |

150 ^a Boltzmann function-based regression analysis results assuming the reported cumulative number of deaths are
151 precise and have uncertainty (a standard deviation of 2.5%), respectively (for detail, refer to the Methods section
152 in the supporting information file and **Figs. 1E, S1**). The same uncertainty was set for Richards function-based
153 regression analysis (for detail, refer to the Methods section and **Fig. S2**)

154
155

Figure 1



156

157 **Figure 1. Fitting the cumulative number of COVID-19 deaths to Boltzmann function**

158 (A, B, C) Boltzmann function-based regressions analysis results on the cumulative numbers of confirmed
 159 COVID-19 cases (panel A) and deaths (panels B and C) in the indicated geographic regions. Parameters
 160 of the established functions for Wuhan City (panels A and B) and for other cities in Hubei (panel C) are
 161 shown in insets. Note: the reported cumulative number of confirmed cases of Hubei Province and Wuhan
 162 City were re-adjusted for data fitting due to the suddenly added cases determined using clinical features
 163 (for details, refer to **Table S1**).

(D) Boltzmann function-based analysis results on the cumulative numbers

164 of 2003 SARS deaths in the indicated regions. Parameters of the established function for mainland China
165 are shown in insets. (E). Regressions analysis results on COVID-19 deaths in Wuhan City by the
166 Boltzmann functions assuming that the relative uncertainty of the data follows a single-sided normal
167 distribution with a mean of 1.0 and a standard deviation of 2.5%. Original data are shown as circles;
168 simulated results are presented as colored lines as indicated. Inserts show key statistics. Results for other
169 regions are presented in **Figs. S1**. (F). Prediction of COVID deaths in Wuhan City by Boltzmann function-
170 based analyses. The real data from Jan 21 to different ending dates were arbitrarily analyzed (colored lines),
171 with the potential total numbers of deaths under these analyses being shown in insets.